The CMG Helicase Bypasses DNA-Protein Cross-Links to Facilitate Their Repair

Graphical Abstract



Authors

Justin L. Sparks, Gheorghe Chistol, Alan O. Gao, ..., Matthias Mann, Julien P. Duxin, Johannes C. Walter

Correspondence

johannes_walter@hms.harvard.edu

In Brief

A collaboration between helicases ensures that DNA damage in the form of DNA-protein cross-links does not derail replication, helping to preserve genome stability.

Highlights

- Replicative helicase CMG bypasses DNA-protein crosslinks on its translocation strand
- RTEL1, an accessory helicase, facilitates bypass by unwinding DNA beyond the DPC
- CMG slows after bypass, likely due to uncoupling from DNA synthesis
- Bypass triggers DPC proteolysis and prevents accidental degradation of CMG



The CMG Helicase Bypasses DNA-Protein Cross-Links to Facilitate Their Repair

Justin L. Sparks,^{1,6} Gheorghe Chistol,^{1,6} Alan O. Gao,^{1,2} Markus Räschle,³ Nicolai B. Larsen,² Matthias Mann,^{2,4} Julien P. Duxin,² and Johannes C. Walter^{1,5,7,*}

¹Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, MA 02115, USA

²Novo Nordisk Foundation Center for Protein Research, Faculty of Health and Medical Sciences, University of Copenhagen, 2200 Copenhagen, Denmark

³Department of Molecular Biotechnology and Systems Biology, Technical University of Kaiserslautern, 67653 Kaiserslautern, Germany ⁴Department of Proteomics and Signal Transduction, Max Planck Institute of Biochemistry, 82152 Martinsried, Germany

⁵Howard Hughes Medical Institute, Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, MA 02115, USA

⁶These authors contributed equally

7Lead Contact

*Correspondence: johannes_walter@hms.harvard.edu

https://doi.org/10.1016/j.cell.2018.10.053

SUMMARY

Covalent DNA-protein cross-links (DPCs) impede replication fork progression and threaten genome integrity. Using Xenopus egg extracts, we previously showed that replication fork collision with DPCs causes their proteolysis, followed by translesion DNA synthesis. We show here that when DPC proteolysis is blocked, the replicative DNA helicase CMG (CDC45, MCM2-7, GINS), which travels on the leading strand template, bypasses an intact leading strand DPC. Single-molecule imaging reveals that GINS does not dissociate from CMG during bypass and that CMG slows dramatically after bypass, likely due to uncoupling from the stalled leading strand. The DNA helicase RTEL1 facilitates bypass, apparently by generating single-stranded DNA beyond the DPC. The absence of RTEL1 impairs DPC proteolysis, suggesting that CMG must bypass the DPC to enable proteolysis. Our results suggest a mechanism that prevents inadvertent CMG destruction by DPC proteases, and they reveal CMG's remarkable capacity to overcome obstacles on its translocation strand.

INTRODUCTION

DNA replication forks encounter many obstacles that challenge genome duplication. Discrete DNA lesions (e.g., pyrimidine dimers) stall the replicative DNA polymerase but not the helicase, leading to helicase-polymerase uncoupling (Byun et al., 2005; Taylor and Yeeles, 2018). Bulkier obstacles block the entire replisome, including the helicase. These include DNA interstrand cross-links (ICLs) (Fu et al., 2011) and DNA-protein cross-links (DPCs) (Duxin et al., 2014). DPCs are formed by ultraviolet light, various chemotherapeutics (e.g., cisplatin), and by endogenous agents such as formaldehyde and topoisomerases (Ide et al., 2011; Stingele et al., 2017). Non-covalent nucleoprotein complexes (e.g., RNA polymerases and tightly bound transcription factors) also interfere with replication fork progression.

Our understanding of DPC repair is evolving rapidly. Early studies showed that DPCs smaller than ${\sim}8$ kDa are excised by nucleotide excision repair whereas larger DPCs require more complex pathways (Ide et al., 2011). In 2014, yeast Wss1 was identified as a DNA-dependent protease that degrades DPCs in S phase (Stingele et al., 2014). Contemporaneous experiments in Xenopus egg extracts showed that when a replication fork collides with a DPC on the leading (DPC^{Lead}) or lagging strand template (DPC^{Lag}), an unknown protease degrades the DPC to a short peptide adduct that is bypassed by translesion DNA polymerases (Figure 1A) (Duxin et al., 2014). Therefore, a proteasemediated DPC repair pathway exists that does not involve a double-strand break intermediate or recombination. Like Wss1, its vertebrate ortholog SPRTN (also named Spartan or DVC1) is a DNA-dependent protease (Lopez-Mosqueda et al., 2016; Stingele et al., 2016; Vaz et al., 2016). The protease activity of SPRTN suppresses genome instability by removing DPCs that block replication fork progression (Lessel et al., 2014; Maskey et al., 2014; Vaz et al., 2016), and SPRTN mutations cause Ruijs-Aalfs syndrome (RJALS), a human genetic disease characterized by genome instability, premature aging, and early onset liver cancer (Lessel et al., 2014). In frog egg extracts, SPRTN and the proteasome have overlapping functions in DPC proteolysis (Larsen et al., 2018). In this setting, SPRTN activity requires that a nascent strand be extended to within a few nucleotides of the DPC, whereas proteasome activity requires DPC ubiquitylation and the presence of single-stranded DNA (ssDNA) near the adduct. How a DPC is selectively degraded without concomitant destruction of the neighboring, stalled replisome is unknown.

Replicative helicases form hexameric rings that unwind DNA by translocating along single stranded DNA ("steric exclusion") (O'Donnell and Li, 2018). Understanding how these essential motors overcome obstacles is an important question. The bacterial DnaB helicase, which translocates 5' to 3' along the lagging strand template, stalls at non-covalent nucleoprotein complexes.



Figure 1. Disappearance of the CMG Footprint at DPC^{Lead} Is Unaffected by Proteolysis or p97 Inhibition

(A) Previous model of replication-coupled DPC repair (Duxin et al., 2014).

(B) Schematic of what happens in the presence of Ub-VS (Duxin et al., 2014).

(C) pDPC^{Lead} or pmeDPC^{Lead} were pre-bound with LacR to prevent one replication fork from reaching the DPC. The plasmids were replicated in mock-depleted or SPRTN-depleted egg extract containing $^{32}P[\alpha]$ -dATP and supplemented with buffer or the p97 inhibitor NMS-873 (p97i). At different times, DNA was recovered and digested with AatII and FspI, separated on a denaturing polyacrylamide geI, and visualized by autoradiography. Gray inset: schematic of nascent leading strand products released by AatII and FspI digestion of pmeDPC^{Lead} or pDPC^{Lead}. The lower autoradiogram shows nascent leading strands generated by the rightward replication fork, and the upper autoradiogram shows both extension products. Blue bracket, CMG footprint (-30 to -37); orange bracket, products stalled at the adducted base (-1 to +1). The percentage of leading strands that approached from the -30 cluster to the -1 cluster was quantified (see STAR Methods), and the mean of n = 5 experiments is graphed. Error bars represent the SD. See Figure S1E for description of -1 to +12 products in lanes 7-12 and 19-24.

(D) pmeDPC^{2xLead} was replicated in SPRTN-depleted egg extracts and supplemented with buffer or p97i. At different times, plasmid-associated proteins were recovered and blotted with the indicated antibodies. Samples were also examined for DPC proteolysis (Figure S1D). A model of CMG unloading from this template is shown in Figure S1G.

To overcome these obstacles, it employs two accessory helicases, UvrD and Rep, which act redundantly (Guy et al., 2009). As 3' to 5' helicases, Rep and UvrD assemble on the leading strand template and cooperate with DnaB in overcoming obstacles. Unlike DnaB, the eukaryotic replicative DNA helicase, CMG (a complex of CDC45, MCM2-7, and GINS), encircles and translocates 3' to 5' along the leading strand template (Fu et al., 2011). While isolated CMG can bypass a biotin-streptavidin (SA) complex on the lagging strand template with the assistance of MCM10, it cannot overcome a leading strand biotin-SA complex (Langston et al., 2017). Interestingly, the large T antigen DNA helicase can bypass a DPC on the translocation strand, perhaps via

transient ring-opening (Yardimci et al., 2012b). Whether CMG progression past nucleoprotein complexes is assisted by accessory helicases is unknown. However, consistent with this idea, in the absence of the 5' to 3' helicases Rrm3 and Pif1, replisomes stall at specific locations in yeast cells (lvessa et al., 2000). These results suggest that, like bacteria, yeast cells overcome obstacles by engaging an accessory DNA helicase that moves on the strand opposite the one hosting CMG.

Based on experiments in *Xenopus* egg extracts, we previously proposed that DPC proteolysis allows the CMG helicase to move past a leading strand DPC (Figure 1A) (Duxin et al., 2014). We now show that CMG can bypass DPC^{Lead} in the absence of DPC proteolysis. Single-molecule imaging demonstrates that GINS does not dissociate during bypass, and CMG slows dramatically after bypass, likely due to temporary uncoupling from the leading strand, which stalls at the DPC. The 5' to 3' helicase RTEL1 facilitates CMG bypass of the DPC, and bypass is required for efficient DPC proteolysis. RTEL1 also promotes fork progression through non-covalent DNA-protein complexes. Our results require a fundamental revision of the events underlying DPC repair and reveal CMG's remarkable capacity to overcome nucleoprotein obstacles.

RESULTS

Leading Strands Rapidly Approach a Non-degradable DPC

To study the mechanism of DPC repair, we covalently trapped the 45 kDa DNA methyltransferase M.Hpall on its recognition site (CCGG) in a circular plasmid (Figure S1A) (Duxin et al., 2014). When the resulting plasmid (pDPC) is replicated in Xenopus egg extracts, leading strands stall 30 nucleotides (nt) from the adducted base due to the footprint of the CMG helicase, which translocates along the leading strand template (Figure 1A) (Duxin et al., 2014). After a pause, synthesis resumes and nascent leading strands approach to within 1 nucleotide of the DPC ("approach"), followed by translesion DNA synthesis (Figure 1A) (Duxin et al., 2014). Because degradation of the DPC roughly coincides with approach, the results suggested that degradation of the DPC allows CMG to move beyond the adduct (Duxin et al., 2014) (Figure 1A). Surprisingly, however, when DPC proteolysis was blocked via ubiquitin depletion, the nascent leading strand still approached the adduct, albeit very inefficiently, suggesting that CMG eventually vacates its position in front of the DPC (Figure 1B) (Duxin et al., 2014). Moreover, the nascent lagging strand was ultimately extended beyond the DPC. Therefore, the replication fork can eventually move past an intact DPC, but the fate of CMG in this process was unclear (Figure 1B).

Given the many roles of ubiquitin signaling in the DNA damage response, we wanted to block degradation of the DPC without ubiquitin depletion and examine the effect on approach. In egg extracts, the DPC is degraded redundantly by SPRTN and the proteasome (Figure S1B) (Larsen et al., 2018). To inhibit SPRTN, we immunodepleted SPRTN from extracts (Figure S1C); to inhibit the proteasome pathway, we used a DPC whose lysine residues were chemically methylated to prevent ubiquitylation (meDPC) (Figure 1C, spheres with bold outlines). Under these conditions, the DPC was not degraded, and we refer to this as

a "stable" DPC (Figure S1D) (Larsen et al., 2018). To ensure that a single fork encountered the stable DPC on the leading strand template ("DPC^{Lead}"), we flanked the meDPC on the right with 48 tandem copies of the Lac repressor (LacR), which blocks arrival of the leftward, converging fork (Figure 1C, gray inset) (Duxin et al., 2014). To monitor progress of the rightward leading strand, we digested the DNA with AatII and Fspl (Figure 1C, gray inset). Strikingly, leading strands approached the DPC with the same kinetics whether or not proteolysis occurred (Figure 1C; compare lanes 1-6 and 7-12, lower autoradiogram; see graph for quantification). The staggered cuts made by Aatll also distinguish leading and lagging strand extension products (Figure 1C, gray inset, pink and purple arrows), which revealed that the lagging strand was readily extended past the stable DPC (Figure 1C, lanes 7-12, top autoradiogram). Leading strand extension past DPC^{Lead} was much slower than lagging strand extension (Figure 1C, lanes 1-6), and the former was further delayed when the DPC was stabilized (Figure 1C, lanes 7-12). Leading strand extension past the stable DPC^{Lead} was strongly inhibited by Rev1 depletion, demonstrating a requirement for TLS (Figure S1E). Collectively, these results show that when ubiquitin levels are normal, the CMG footprint disappears from the DPC, and the lagging strand is extended past the DPC with the same kinetics whether or not the DPC undergoes proteolysis.

CMG Bypasses a Stable DPC

The disappearance of the CMG footprint at a stable DPC could be due to CMG dissociation from the DPC (in which case another helicase enables lagging strand extension past the DPC) (Figure S1Fi) or CMG bypass of the DPC (Figure S1Fii). To distinguish between these possibilities, we examined approach in the presence of an inhibitor of the p97 ATPase (p97i), which extracts CMG from chromatin during replication termination and ICL repair (Fullbright et al., 2016; Maric et al., 2014; Moreno et al., 2014). If CMG has to dissociate from the stalled replisome to enable approach, p97i should delay approach. As shown in Figure 1C, p97i did not slow the kinetics of leading strand approach to or lagging strand extension past a stable DPC, suggesting that CMG bypassed the DPC. We verified that p97i prevented CMG unloading from a DPC-containing plasmid (Figure 1D), and we infer that such unloading normally occurs after DPC bypass (Figure S1G). DPC bypass did not involve new origin firing (Figure S1H), nor did it require the FANCM ATPase (Figures S1I and S1J) or ATR signaling (Figure S1K), suggesting bypass is mechanistically distinct from ICL traverse (Huang et al., 2013; Ling et al., 2016). Together, our data strongly suggest that CMG bypasses a bulky DPC on the translocation strand.

ssDNA Downstream of a Stable DPC Facilitates CMG Bypass

We speculated that a 5' to 3' accessory helicase loads onto the lagging strand template and unwinds DNA past the DPC^{Lead} to assist CMG bypass (Figure S2i). To explore this idea, we placed a second meDPC on the lagging strand template 15 nucleotides upstream of meDPC^{Lead} (pmeDPC^{Lag/Lead}), as this should block unwinding past the DPC by a 5' to 3' helicase (Figure S2ii). For comparison, we placed the second DPC 15 nucleotides downstream of the leading strand DPC (pmeDPC^{Lead/Lead}) as this



The indicated plasmids were pre-incubated with LacR, replicated in SPRTN-depleted egg extract, and analyzed as in Figure S1J. Approach was used as a proxy for CMG bypass and quantified as in Figure 1C (see STAR Methods). Figure S2 depicts the proposed events on each plasmid. The mean of n = 3 experiments is graphed. Error bars represent the SD.

should allow some DNA unwinding beyond the DPC (Figure S2iii). Consistent with our hypothesis, CMG bypass was severely inhibited by the upstream but not the downstream lagging strand DPC (Figure 2, lanes 19–24 versus 25–30; graph for quantification). Strikingly, a 40 nucleotide ssDNA bubble placed downstream of the DPC fully rescued bypass in the presence of the upstream DPC^{Lag}, further suggesting that DNA unwinding past the DPC is critical for bypass (Figures 2, lanes 31–36, and S2iv). Remarkably, bypass was now even faster than on the DNA template lacking any lagging strand obstruction (Figure 2, compare lanes 31–36 with lanes 7–12). Interestingly, two tandem meDPCs placed 15 nucleotides apart on the leading strand template also severely impaired bypass (Figure 2, lanes 13–18), suggesting that a single CMG cannot simultaneously accommodate two DPCs during bypass. Together, the data suggest that generation of ssDNA downstream of DPC^{Lead} is rate-limiting for CMG bypass, consistent with assistance by a 5' to 3' helicase.

RTEL1 Promotes Efficient CMG Bypass of a Stable DPC

Vertebrates contain six 5' to 3' DNA helicases (RTEL1, FANCJ, PIF1, DDX3, DDX11, and XPD). Using mass spectrometry (PP-MS), we detected all of these helicases except DDX11 on



(legend on next page)

chromatin in egg extracts (Figure 3A) (Larsen et al., 2018), and three (PIF1, RTEL1, and FANCJ) bound the plasmid in a replication-dependent fashion. Depletion of PIF1, FANCJ, or DDX3 had no significant effects on bypass of a stable DPC^{Lead} (data not shown; we did not examine XPD or DDX11). In contrast, immunodepletion of RTEL1 (Figures 3B and S3A for explanation of RTEL1 isoforms in lane 1 of Figure 3B) caused leading strands to initially stall at a greater distance from the stable DPC^{Lead} (Figure 3C, -38 to -44 cluster) and greatly delayed their approach to the adduct (Figure 3C; graph for quantification). Furthermore, RTEL1 depletion delayed lagging strand advance beyond DPC^{Lead} (Figure 3D, lanes 6–10, upper autoradiogram). These results indicate that RTEL1 is required for efficient CMG bypass and that bypass allows new Okazaki fragment priming downstream of the adduct. RTEL1 depletion had no effect on the efficiency of DNA synthesis (Figure S3B). CMG bypass was rescued by wild-type recombinant RTEL1 but not an ATPase deficient RTEL1 mutant (RTEL1K48R), which further inhibited bypass (Figures 3C and S3C). RTEL1 depletion had only a modest effect on bypass when a single-stranded DNA bubble was placed downstream of the DPC, and it did not further inhibit bypass when meDPC^{Lag} was present upstream (Figure S3D). Our data indicate that RTEL1 enables efficient CMG bypass of intact DPC^{Lead} by generating ssDNA beyond the adduct. The bypass observed in RTEL1-depleted extracts (Figure 3C) could be due to incomplete RTEL1 depletion, the presence of partially compensating helicases, or helicase-independent bypass. The bypass defect observed when a single fork collided with DPC^{Lead} in RTEL1depleted egg extract was rescued by IPTG, which disrupted the LacR array and allowed a second fork to converge on the DPC (Figure 3E, lanes 18-22 versus 23-28; graph for quantification). This rescue was not due to CMG unloading as it still occurred in the presence of p97i. We conclude that DPC bypass requires ssDNA beyond the adduct, which can be created by an accessory helicase or a converging replication fork.

M.Hpall likely interacts intimately with both strands of the double helix (Klimasauskas et al., 1994). Therefore, covalent coupling of M.Hpall to one strand is expected to hyperstabilize the underlying DNA duplex. To test whether RTEL1 disrupts such a hyperstable duplex, we examined DPC^{Lag}, in which M.Hpall should stabilize the underlying DNA without blocking the translocation strand. As shown in Figure 3D, RTEL1 depletion delayed bypass of DPC^{Lag}, but the delay was less pronounced than at DPC^{Lead} (Figures 3D and S3E). Collectively, these obser-

vations suggest that RTEL1 not only helps CMG disrupt hyperstable DNA duplex but also to overcome obstacles that are covalently attached to CMG's translocation strand.

Efficient DPC Proteolysis Requires RTEL1

The kinetics of CMG bypass were identical whether or not the DPC was degraded (Figure 1C). This observation strongly implies that CMG normally bypasses the DPC before proteolysis, and it raised the question of whether bypass might be a pre-requisite for proteolysis. To address this, we replicated a plasmid containing non-methylated DPC^{Lead} in extract containing SPRTN so that both proteolysis pathways were active. The extract was mock-depleted or depleted of RTEL1. At different times, we isolated the plasmid under stringent conditions so that only covalently attached M.Hpall was recovered. After DNA digestion, immunoblotting revealed M.Hpall polyubiquitylation, followed by a decline in M.Hpall levels, which reflects replication-dependent proteolysis (Figure 4A, lanes 1-6) (Larsen et al., 2018). Strikingly, M.Hpall degradation was delayed 15-20 min in extracts depleted of RTEL1 (Figure 4A). This defect was rescued by wild-type RTEL1 but not the ATPase mutant. RTEL1 was required for efficient CMG bypass even when the DPC could be degraded, consistent with CMG bypass normally preceding DPC proteolysis (Figure 4B). In the absence of RTEL1, M.Hpall proteolysis was delayed but not eliminated (Figure 4A), which is explained by the substantial CMG bypass that still occurred in these conditions (Figure 4B). Together, the data indicate that RTEL1-mediated unwinding past the DPC is essential for its efficient proteolysis.

RTEL1-Dependent DPC Bypass Promotes SPRTN Activity

To examine whether RTEL1 is required for SPRTN activity, we examined meDPC, which is not susceptible to the proteasome but can be degraded by SPRTN (Larsen et al., 2018). The action of SPRTN was visible from the appearance of a specific M.Hpall degradation fragment that was absent in Δ SPRTN extract (Figure 4C) (Larsen et al., 2018). In the absence of RTEL1, accumulation of the SPRTN-specific fragment was delayed (Figure 4C, compare lanes 1–5 and 11–15). As seen for DPC bypass, these defects were rescued by wild-type but not ATPase-deficient RTEL1, indicating that DNA unwinding past the DPC by RTEL1 is required for SPRTN activity. To address whether CMG bypass itself is required, we examined the effect of tandem leading strand

Figure 3. RTEL1 Is Required for Efficient CMG Bypass

⁽A) Recovery of 5' to 3' helicases in the mass spectrometry dataset of Larsen et al. (2018). Relative abundance of each protein in the specified conditions is expressed as a Z score with yellow indicating higher abundance. Where indicated, Geminin was added to block replication initiation.

⁽B) Mock-depleted, RTEL1-depleted, and RTEL1-depleted egg extracts supplemented with wild-type RTEL1 or RTEL1-K48R were blotted with RTEL1 and ORC2 (loading control) antibodies.

⁽C) pmeDPC^{Lead} was replicated in the indicated extracts, and supplemented with buffer, wild-type RTEL1, or RTEL1-K48R. Leading strand approach was visualized as in Figure S1J and quantified as in Figure 1C. The mean of n = 3 independent experiments is graphed. Error bars represent the SD.

⁽D) pmeDPC^{Lead} or pmeDPC^{Lag} was replicated in the indicated extracts and analyzed as in Figure 1C. CMG bypass was quantified as in Figure 1C and is shown in Figure S3E.

⁽E) pmeDPC was replicated in the indicated extracts and supplemented with IPTG (at 5 min after replication initiation) and/or p97i, as indicated. Leading strand approach was visualized as in Figure S1J and quantified as in Figure 1C. The mean of n = 3 independent experiments is graphed. Error bars represent the SD. The slower CMG bypass observed in RTEL1-depleted extract containing IPTG relative to mock-depleted extract (light blue versus red graph) is largely accounted for by the fact that CMG progression through the lacO array is delayed by LacR, even in the presence of IPTG, as seen from the slower appearance of resolved, linear species in Figure S3F (compare light blue versus red graphs).



(legend on next page)

meDPCs, which severely impair CMG bypass (Figure 2), presumably without affecting RTEL1's ability to unwind past the adducts. As shown in Figure 4D, tandem meDPCs severely inhibited the appearance of the SPRTN-specific M.Hpall fragment compared to single meDPCs, which allow CMG bypass. Our data indicate that RTEL1-dependent DNA unwinding and CMG bypass of the DPC are both required for the SPRTN pathway.

We next addressed whether RTEL1 affects the proteasome pathway. A time course revealed that in the absence of RTEL1, the appearance of highly ubiquitylated M.Hpall species was delayed, suggesting that RTEL1 is required for efficient DPC ubiquitylation (Figure 4E). To specifically examine the effect of RTEL1 on the proteasome pathway, we replicated pDPC^{2xLeads} in SPRTN depleted-extract with or without RTEL1 depletion and examined DPC proteolysis, using MG262 addition as a positive control for proteasome inhibition. As shown in Figure 4F. RTEL1 depletion stabilized M.Hpall to a similar extent as MG262 in ΔSPRTN extract (compare lanes 11-15 with 16-20), consistent with RTEL1 functioning in the proteasome pathway. Finally, in RTEL1-depleted extracts, chromatin-binding of SPRTN was reduced, and binding of the PSA3 proteasome subunit was delayed (Figure S4A). Together, our experiments indicate that RTEL1 is required for both proteolysis pathways.

We wanted to know whether RTEL1 regulates DPC proteolysis independently of promoting CMG bypass. We therefore employed a substrate containing a ssDNA gap across from the DPC, in which DPC proteolysis by SPRTN and the proteasome occurs in the absence of a replication fork (Figure S4C, lanes 1–6) (Larsen et al., 2018). Importantly, in this replication-independent setting, RTEL1 depletion had no effect on DPC ubiquitylation and proteolysis (Figure S4C) or production of the SPRTN-specific M.Hpall degradation fragment (Figure S4D). These results suggest that RTEL1 does not directly regulate SPRTN or the proteasome but does so indirectly, by stimulating CMG bypass.

Direct Observation of DPC Bypass by Single-Molecule Analysis

To investigate CMG dynamics during DPC repair, we developed a single-molecule assay called KEHRMIT (kinetics of the eukaryotic helicase by real-time molecular imaging and tracking) that is similar to an approach developed in yeast extracts (Duzdevich et al., 2015). DNA replication was initiated on stretched λ DNA (Figures 5Ai–5Aiii) using GINS-depleted egg extract reconstituted with active, recombinant GINS labeled on its Psf3 subunit with Alexa Fluor 647 (Figures 5B and S5A–S5D; rGINS^{AF647} labeling efficiency ≥90%). After ~2 min, when only a few origins

per λ DNA had fired, we flowed in GINS-depleted extract lacking rGINS^{AF647} to remove background fluorescence and prevent further origin firing (Figure 5Aiv). Subsequent imaging of AF647 (Figure 5Av) revealed that each replication origin gave rise to two diffraction-limited spots of comparable intensity that traveled in opposite directions (Figure 5C, green). Spots photobleached in a single step, indicating they contain a single rGIN-SAF647 molecule (Figure S5E). CMGs traveled at an average velocity of ~400 nucleotides per min (Figure 5D), consistent with fork rates on immobilized DNA templates (Loveland et al., 2012). Furthermore, rGINSAF647 migrated at the leading edge of nascent DNA tracts (Figure 5C, blue; see legend), indicating that each AF647 spot represents CMG at a replication fork. Labeled CMGs traveled on average 5.3 kb before disappearing (Figure 5E). For molecules where the DNA remained attached until the end of the experiment, the nascent DNA tract continued to grow even after loss of AF647 signal (Figure S5F). Given that there is no free pool of GINS during the imaging phase (Figures 5Aiv and 5Av), the loss of AF647 signal reflects photobleaching and not GINS exchange. We conclude that GINS is highly processive and that KEHRMIT is a powerful means to examine CMG dynamics during replication.

To examine the mechanism of DPC repair, we immobilized linear DNA containing two site-specific DPCs labeled at their C termini with Alexa Fluor 568 (AF568) (Figure 6A). If an origin fires between the two lesions, both helicases encounter a DPC^{Lead} (Figure 6B, top); otherwise, the inward moving CMG first encounters DPC^{Lag} and then DPC^{Lead} (Figure 6B, bottom). We first conducted KEHRMIT on meDPC in SPRTN-depleted extract to inhibit DPC proteolysis and thereby maximize the possibility of observing bypass. Strikingly, we saw many instances in which CMG paused at DPC^{Lead} and then underwent bypass (Figure 6C). Interestingly, CMG slowed down dramatically after bypass, as described below. To quantify bypass events and determine their timing, we applied the following, stringent criteria: (1) both CMG^{AF647} and DPC^{AF568} signals were present before, during, and after bypass; (2) CMG traveled at least 1 pixel (~500 nucleotides) away from the DPC after bypass; and (3) CMG translocated for at least 3 time points (3 min) before and after bypass. Based on this algorithm, 42%-56% of CMGs that encountered meDPC^{Lead} in SPRTN-depleted extract unambiguously bypassed the intact DPC ("BID" events) (Figures 6D and S6A), and CMG paused at meDPC^{Lead} for 15 min on average before undergoing bypass (Figure 6E). We observed four other classes of events (Figures 6D, see legend for a detailed description, and S6B for examples). Many of these non-BID events

Figure 4. RTEL1 Is Required for Efficient DPC Proteolysis

⁽A) $pDPC^{2xLead}$ was replicated in the indicated extracts and supplemented with buffer, wild-type RTEL1, or RTEL1-K48R. Plasmid was recovered under stringent conditions, the DNA digested, and the resulting samples blotted for Hpall. Signal from the entire lane was quantified, and peak signal was assigned a value of 100%. The mean of n = 3 independent experiments is graphed. Error bars represent SD.

⁽B) Parallel reactions to those in (A) were supplemented with $[\alpha$ -³²P]-dATP. Leading strand approach was visualized as in Figure S1J and quantified as in Figure 1C. The mean of n = 3 independent experiments is graphed. Error bars represent SD.

⁽C) pmeDPC^{2xLead} was replicated in the indicated extracts. Samples were processed by the pull-down procedure described in (A). Short and long (lower panel) exposures of the same blot are shown.

 ⁽D) pmeDPC^{2xLead} or pmeDPC^{2xLead} were replicated in non-depleted extract. Stringent plasmid pull downs were performed as in (A) and presented as in (C).
(E) pDPC^{2xLead} was replicated in the indicated extracts, and stringent plasmid pull-down was performed as in (A). RTEL1 depletion was verified in Figure S4B.
(F) pDPC^{2xLead} was replicated in the indicated egg extracts that also contained DMSO or MG262. Stringent plasmid pull-down was performed as in (A).



probably involve meDPC^{Lead} bypass that could not be detected due to the slow rate of CMG progression after bypass, the stringent criteria for BID events, and the premature loss of CMG or DPC signal due to photobleaching (Figure S6C) or DNA breakage. Therefore, the actual DPC^{Lead} bypass efficiency was probably substantially higher than 50%.

When CMG encountered meDPC^{Lag} in SPRTN-depleted egg extract, it paused on average for only 3 min before moving past the adduct (Figures 6E and 6F), consistent with ensemble analysis of leading strands (Duxin et al., 2014). In 44%–49% of cases, CMG unambiguously bypassed meDPC^{Lag} (Figure S6D, BID). In 47%–51% of meDPC^{Lag} encounters, meDPC^{Lag} became mobile after helicase bypass and then tracked with CMG (Figures S6D, B+M, and S6E). Because the DNA template is immo-

Figure 5. KEHRMIT: A Single-Molecule Assay for CMG Dynamics

(A) Schematic of KEHRMIT assay.

(B) Coomassie-stained SDS-PAGE gel of recombinant GINS before and after sortase labeling of Psf3 with AF647, which shifts its mobility (arrow).
(C) Kymogram of a replication bubble from a KEHRMIT experiment. Green, GINS^{AF647} signal. Blue, Fen1^{mKikGR}, a fluorescent protein that binds nascent DNA (Loveland et al., 2012).

(D and E) Beeswarm plots of CMG speed (D) and CMG processivity (E) (i.e., distance traveled) measured via KEHRMIT (dots represent n = 218 individual helicase molecules). Blue line, mean; gray box, 95% confidence interval (CI) estimated by bootstrapping.

bilized via 3'-biotins, meDPC^{Lag} probably becomes mobile when the outward moving replisomes reaches the end of the template, which liberates and allows chromatin compaction of the sister chromatid containing the bypassed DPC (Figure S6F). Taking into account BID and B+M events, meDPC^{Lag} bypass efficiency exceeded 95% (Figure S6D). In conclusion, KEHRMIT shows that CMG efficiently bypasses stable DPC^{Lead} and DPC^{Lag}, and this process does not involve GINS dissociation.

To address whether CMG bypasses DPC^{Lead} when proteolysis is not impaired, we examined a non-methylated DPC in extract containing SPRTN. As expected, signatures involving early loss of DPC^{Lead} increased when proteolysis was allowed (Figure 6D, DD⁻ and DD⁺). In this setting, 11%–14% of CMGs exhibited unambiguous DPC^{Lead} bypass (Figure 6D, BID). We suspect that many DD⁺ events represent bona-fide bypass, but failed to meet the BID criteria because DPC proteolysis occurred before CMG traveled far enough beyond the DPC. In support of this inter-

pretation, the probabilities of BID and DD⁺ events add up to 40%–45% for degradable DPC^{Lead} – similar to the likelihood of unambiguous bypass (BID 42%–56%) for stabilized DPC^{Lead} (Figure 6D). As noted above, any bypass events in the DT, DD⁻, and CD categories go undetected due to the premature disappearance of DPC and/or CMG signal. Thus, 11%–14% is likely a gross underestimate of the true efficiency with which CMG bypassed DPC^{Lead}. In conclusion, our results are consistent with the model that CMG bypasses DPC^{Lead} prior to proteolytic processing.

Differential Kinetics of DPC^{Lead} versus DPC^{Lag} Proteolysis

In these experiments, the AF568 fluorescence signal disappears when the C terminus of Hpall is degraded, providing a



Figure 6. Direct Observation of DPC Bypass by CMG

(A) Stretched and immobilized DNA-DPC substrate. DNA was stained with sytox orange (top). The DPC was labeled on its C terminus with AF568 (middle). Bottom: merge.

(B) Cartoons depicting how the location of initiation determines whether CMGs encounter DPC^{Lead} (top) or DPC^{Lag} followed by DPC^{Lead} (bottom).

(C) Kymogram of a meDPC substrate undergoing replication in SPRTN-depleted extract from an origin that fired between the DPCs. Both CMGs undergo DPC bypass. Images were acquired at 1 frame/min. Green, AF647; pink, AF568.

(D) Quantification of five different classes of $CMG-DPC^{Lead}$ encounters in n = 2 independent biological repeats: (1) BID (bypass of intact DPC), representing unambiguous bypass events; (2) DD+ (DPC disappeared first), followed by CMG departure from the pause site; when proteolysis was inhibited (meDPC, Δ SPRTN extract), DD⁺ events likely involve DPC bypass but do not meet the BID criteria due to DPC photobleaching; (3) DD⁻ (DPC disappeared first), without CMG departure from the pause site, including potential bypass events where CMG photobleached or DNA ruptured soon after the DPC signal vanished; (4) CD (CMG disappeared first), likely due to photobleaching, obscuring any subsequent bypass events; and (5) DT (CMG and DPC disappeared together), including events where the experiment ended or the DNA ruptured before bypass could be detected. Due to rounding errors, probabilities may not add up to 100%. n represents the number of molecules.

(E) Beeswarm plot of the time needed to bypass meDPC^{Lead} or meDPC^{Lag} in SPRTN-depleted extract. Blue lines and gray boxes correspond to the mean and the 95% CI for the mean, respectively. n represents the number of molecules.

(F) Same as (C) but showing a kymogram in which an origin fired to the right of both lesions. The leftward-moving helicase (green) first bypassed a meDPC^{Lag} in ~1 min, then reached a meDPC^{Lead} where it paused.

(G) Beeswarm plot of DPC lifetime after CMG arrival at the lesion. Blue lines, gray boxes, and n as in (E).

(H) Beeswarm plot of CMG speed during approach to (Appr.) and departure from (Dep.) DPC lesions versus the speed of aphidicolin-uncoupled helicases. Blue lines, gray boxes, and n as in (E). The aphidicolin condition was performed on λ DNA.

(I) Kymogram of CMG-meDPC^{Lead} encounter (SPRTN-depleted extract) that resulted in DPC bypass and CMG uncoupling, followed by apparent re-coupling (white arrow).

single-molecule measurement of DPC proteolysis. In most instances, the loss of the AF568 signal was due to DNA replication and not photobleaching (Figure S6G). Importantly, DPC^{Lag} fluorescence disappeared ~7 min after CMG arrival, whereas DPC^{Lead} disappearance took ~25 min (Figure 6G). This result is consistent with our finding that proteolysis depends on bypass (Figure 4D), and bypass takes much longer for DPC^{Lead} than DPC^{Lag} (Figure 6E). As expected, when the DPC was methylated and examined in SPRTN-depleted extract, both meDPC^{Lead} and meDPC^{Lag} were greatly stabilized (Figure 6G). In conclusion, the difference between DPC^{Lag} and DPC^{Lead} destruction is accounted for by the different kinetics with which these lesions are bypassed, consistent with bypass being required for proteolysis.

CMG Slows Down after DPC^{Lead} Bypass

We investigated how CMG's velocity is affected by collision with DPC^{Lead} and DPC^{Lag}. Strikingly, while CMG continued moving at 400–500 nucleotides/min after bypassing DPC^{Lag}, it slowed to ~80 nucleotides/min after bypassing DPC^{Lead} (Figure 6H). We speculated that CMG slowing was due to its uncoupling from leading strand synthesis, which pauses for an extended time at



Figure 7. RTEL1 Is Required for CMG Bypass of Non-covalent Nucleoprotein Complexes

(A) Top: DNA structures generated with and without XmnI digestion before and after forks progress through the LacR array.

(B) pLacO₃₂ was pre-incubated with LacR and replicated in the indicated egg extracts containing $[\alpha^{-32}P]$ -dATP. DNA was recovered, digested with the single cutter Xmnl, resolved by native agarose gel electrophoresis, and visualized by autoradiography.

(C) DNA samples from (B) were nicked with Nt.BspQl to release the rightward leading strand (red arrow), separated on a denaturing polyacrylamide gel, and visualized by autoradiography. *lacO* sites are located ~30 nucleotides downstream of each stalling product (Dewar et al., 2015).

(legend continued on next page)

DPC^{Lead} but not DPC^{Lag} (Figure 1C) (Duxin et al., 2014). To independently assess how CMG uncoupling from the leading strand impacts helicase velocity, we added aphidicolin, a potent inhibitor of replicative polymerases. In the presence of aphidicolin, CMG slowed dramatically (Figure S6H) to an average speed of only ~55 nucleotides/min (Figure 6H) despite continued high processivity (Figure S6I). The data strongly imply that the slow rate of CMG translocation after DPC^{Lead} bypass is due to CMG uncoupling. In 5% of cases, we observed that after DPC^{Lead} bypass, CMG traveled slowly for several minutes before resuming rapid translocation, indicative of recoupling (Figure 6I). The small number of putative recoupling events is consistent with the low efficiency of TLS after extracts have been depleted. Our results suggest that after CMG bypasses DPC^{Lead}, CMG uncouples from the leading strand and slows down, minimizing ssDNA generation during TLS past the peptide adduct.

RTEL1 Promotes Efficient CMG Progression Past Non-covalent Nucleoprotein Complexes

As shown in Figure 3D, RTEL1 is required to efficiently bypass a lagging strand DPC, probably because it helps unwind the DNA underlying the DPC. If this interpretation is correct, RTEL1 should also be required for replicative bypass of non-covalent nucleoprotein complexes that stabilizes the duplex. To test this prediction, we replicated a plasmid containing an array of 32 lacO sites bound by LacR. At different times, plasmid was recovered and cut with XmnI before native gel electrophoresis (Figure 7A). In this setting, replication forks converge on the outer edges of the LacR array, generating a discrete X-shaped intermediate whose mobility decreases as forks slowly progress through the array (Figure 7B, lanes 1-6) (Dewar et al., 2015). When forks meet, the X-shaped species are converted into linear daughter molecules. As shown in Figure 7B, RTEL1 depletion severely delayed the accumulation of linear molecules, and this effect was partially rescued by RTEL1^{WT} but not RTEL1^{K48R}. To examine fork progression through the LacR array at higher resolution, DNA was nicked near the lacO sites with Nt.BspQI, and the radioactive nascent strands were separated on a urea PAGE gel, which reveals fork pausing \sim 30 nucleotides upstream of each lacO site (Figure 5C, lanes 1-6) (Dewar et al., 2015). Based on this analysis, fork progression through the array was also severely compromised in extracts lacking RTEL1, and the defect was reversed with RTEL1^{WT} but not RTEL1^{K48R} (Figure 7C, red boxes). Replisome progression does not require LacR proteolysis (Figure S7A). We conclude that RTEL1 is required for the eviction of non-covalent nucleoprotein complexes.

DISCUSSION

Our results establish a comprehensive framework to understand how the vertebrate replisome overcomes covalent and noncovalent nucleoprotein obstacles (Figure 7D).

Disruption of Non-covalent Nucleoprotein Complexes by RTEL1

The most common nucleoprotein obstacles encountered by replication forks are non-covalent. We showed that CMG progression past a LacR array requires RTEL1, which translocates 5' to 3' on the lagging strand template. Together with prior work (Guy et al., 2009; Ivessa et al., 2000), a conserved mechanism emerges in which replication forks employ an accessory helicase on the strand opposite the one hosting the replicative DNA helicase. In egg extracts and cells, RTEL1 is constitutively associated with replication forks in the absence of experimentally induced DNA damage (Figure 3A) (Vannier et al., 2013), suggesting RTEL1 travels with the replisome. We propose that RTEL1 unwinds the DNA underlying non-covalent nucleoprotein complexes, thereby disrupting the obstacle and allowing fork progression.

CMG Bypasses DPCs on Both Strands

Less commonly, forks encounter covalent DPCs. We previously showed that the disappearance of the CMG footprint at DPC^{Lead} correlates with the latter's proteolysis, and when DPC^{Lead} proteolysis is blocked by ubiquitin depletion, loss of the footprint is dramatically delayed. On this basis, we proposed that CMG progression past DPC^{Lead} requires DPC proteolysis (Duxin et al., 2014). However, we now show that when ubiquitin levels are normal, CMG can readily bypass a stable DPC^{Lead}. Thus, when DPC proteolysis is blocked by inhibiting the SPRTN and proteasome pathways, the CMG footprint disappears, even in the absence of p97 activity, implying that CMG bypasses the DPC without dissociating. In support of this interpretation, singlemolecule imaging shows that the same molecule of CMG that encounters a DPC^{Lead} also travels past the adduct. Our previous observation that loss of the CMG footprint was inhibited when DPC proteolysis was blocked by a non-specific DUB inhibitor was most likely due to pleiotropic consequences of ubiquitin depletion (Duxin et al., 2014).

We envision two possible mechanisms of DPC^{Lead} bypass. In one, CMG threads the denatured DPC through its central channel (Figure S7B). Because most DPCs should be attached to DNA at an internal amino acid, this mechanism would require CMG's central pore to accommodate ssDNA and two polypeptide chains. Notably, the time required for DPC bypass is not influenced by methylation status (Figure S6J compare meDPC versus DPC; Figure 1C for ensemble data) or when MG262 was added to block proteasome activity (data not shown), suggesting that ubiquitylation of the DPC does not adversely affect bypass. This observation disfavors the "threading" model, which predicts that ubiquitylated DPC would be much more difficult to accommodate in CMG's central channel, even after it is unfolded. In a second model, the MCM2-7 ring opens, allowing CMG to slide past the DPC (Figure S7C). This model is consistent with current evidence that the MCM2-7 ring opens during

⁽D) Model of replisome bypass of nucleoprotein barriers. When the replisome encounters a non-covalent nucleoprotein complex, RTEL1 and CMG cooperate to unwind the DNA underlying the protein, leading to its displacement and immediate resumption of fork progression. At a covalent DPC, RTEL1 translocates along the undamaged lagging strand template, exposing ssDNA that facilitates CMG bypass. Given the stable interaction of pol ε (gray oval) with CMG (Langston et al., 2014), we envision that it bypasses the DPC with CMG. After CMG bypass, the DPC undergoes proteolysis by SPRTN or the proteasome. Finally, the leading strand is extended past the peptide adduct using translesion synthesis polymerases.

replication licensing and initiation (Bochman and Schwacha, 2008; Fu et al., 2011; Samel et al., 2014). Future experiments will be required to distinguish between the above models.

We recently discovered that upon collision of a replication fork with a DPC, the E3 ubiquitin ligase TRAIP promotes DPC ubiquitylation and proteolysis (Larsen et al., 2018). TRAIP-dependent DPC ubiquitylation does not require RTEL1 and therefore occurs independently of CMG bypass (Larsen et al., 2018). Surprisingly, ubiquitylation of the DPC by TRAIP appears to contribute to efficient CMG bypass. Given that DPC methylation, which prevents DPC ubiquitylation, does not affect bypass (Figure 1C), we speculate that methylation mimics the effect of ubiquitylation on bypass. The role of post-translational modifications in DPC bypass is an important area of future investigation.

The requirement for RTEL1 in CMG bypass is largely abolished by a converging fork or when a ssDNA bubble is placed downstream of the DPC, suggesting that the primary function of RTEL1 is to generate ssDNA downstream of the lesion. We favor the idea that ssDNA created beyond the DPC allows the breached CMG to re-engage with DNA beyond the DPC. Given that the non-catalytic N-terminal tier of CMG resides at the leading edge of the fork (Douglas et al., 2018; Georgescu et al., 2017), the N terminus might re-close around ssDNA while the C-terminal ATPase domain is still engaged in bypassing the DPC. Such a mechanism would help avoid accidental dissociation of CMG during DPC bypass.

Although CMG movement past DPC^{Lead} and DPC^{Lag} both constitute "bypass" events, the underlying mechanisms are different. While CMG pauses briefly at DPC^{Lag}, it stalls for an extended period at DPC^{Lead}. In addition, the absence of RTEL1 causes greater CMG stalling at DPC^{Lead} versus DPC^{Lag}. Given CMG's translocation along the leading strand template (Fu et al., 2011), these data indicate that the primary role of RTEL1 at DPC^{Lag} is to assist CMG in disrupting non-covalent interactions between the DPC and the underlying DNA. In contrast, bypassing a leading strand DPC additionally requires RTEL1dependent CMG stepping over or around the covalent DPC-DNA linkage. We previously investigated the effect of biotinstreptavidin roadblocks on CMG translocation. Whereas CMG bypass of these obstacles on the lagging strand template takes only 5-10 min, the same barriers on the leading strand template stall CMG for ~20-40 min (Fu et al., 2011), as seen for DPCs. In summary, although CMG can overcome lagging and leading strand barriers, the latter is more complex as it likely involves profound remodeling of CMG and/or the DPC.

CMG Slows Down after Uncoupling from the Leading Strand

At sites of leading strand DNA damage or when DNA synthesis is inhibited, CMG uncouples from the point of synthesis, leading to ssDNA generation and ATR checkpoint activation (Byun et al., 2005; Taylor and Yeeles, 2018). However, the dynamics of the uncoupled CMG are unknown. We showed that in the presence of aphidicolin, the rate of CMG translocation slows 8-fold. Given the slow DNA unwinding by uncoupled prokaryotic DNA helicases (Graham et al., 2017; Stano et al., 2005), optimal CMG activity likely requires coupling with DNA synthesis. Importantly, after CMG bypass of DPC^{Lead}, the rate of CMG translocation also slowed dramatically, indicative of CMG uncoupling from the leading strand, which stalls at the DPC. Slow translocation of the uncoupled CMG limits the amount of ssDNA generated during replication stress and therefore reduces the likelihood of DNA breakage, accelerates recoupling of the leading strand with CMG after TLS, and may promote template switching by limiting physical separation of the sister chromatids.

CMG Bypass Is Required for DPC Proteolysis

The following evidence indicates that CMG bypass precedes and is essential for DPC proteolysis. First, the CMG footprint disappears at DPCs with the same kinetics whether or not proteolysis occurs (Figure 1C), implying that CMG bypass normally precedes proteolysis. Second, we observe bypass of degradable DPCs in single-molecule experiments. Third, when we inhibit CMG bypass with RTEL1 depletion or tandem DPCs, proteolysis is compromised. Fourth, proteolysis takes longer for DPC^{Lead} than DPC^{Lag}, consistent with the slower kinetics of DPC^{Lead} bypass. Finally, DPC proteolysis by SPRTN requires that the leading strand advance to within a few nucleotides of the DPC (Larsen et al., 2018), which is only possible if CMG has moved out of the way. Altogether, the data strongly support the idea that CMG bypass is a pre-requisite for DPC proteolysis.

The relationship between CMG bypass and the proteasome is less clear. RTEL1 depletion impairs the production of long ubiquitin chains on the DPC, and it mimics MG262 addition in SPRTN-depleted extracts, implicating RTEL1 in the proteasome pathway. However, it is unclear whether RTEL1-dependent DNA unwinding at the DPC is sufficient to trigger proteasome activity, or whether CMG bypass is also required. Consistent with the former possibility, ssDNA is sufficient to trigger proteasomemediated DPC proteolysis in the absence of a leading strand at the lesion (Larsen et al., 2018). A mechanism in which proteasome activity requires RTEL1 unwinding but not CMG bypass would allow the destruction of DPCs that cannot be bypassed (e.g., because they are too large).

The "bypass first" mechanism we describe here is well suited to enhance genome stability. First, CMG movement past DPCs before proteolysis might reduce the probability that the helicase is accidentally destroyed, which is crucial as there is no known pathway to reload MCM2-7 *de novo* in S phase. Second, if CMG bypass of DPC^{Lead} occurs but TLS fails, the lagging strand is still extended past the adduct (Figure 1C; upper autoradiogram). In this case, the leading strand could also be extended past the DPC via template switching (Figure S7D) or re-priming (Taylor and Yeeles, 2018).

RTEL1-Dependent Disruption of Nucleoprotein Complexes *In Vivo*?

RTEL1 regulates telomere homeostasis, and hypomorphic mutations in the human gene cause the telomere shortening Hoyeraal-Hreidarsson syndrome, which is characterized by bone marrow failure, immunodeficiency, and growth retardation (Vannier et al., 2014). RTEL1 resolves telomeric G-quadruplexes and dissolves T-loops, both of which facilitate telomere replication. While homozygous RTEL1 mutations are lethal, deletion of RTEL1's PIP box leads to a general replication fork progression

defect (Vannier et al., 2013). In egg extracts, the PIP box of RTEL1 is not required to bypass DPCs (data not shown), indicating that *Xenopus* RTEL1 can bind to replication forks via other means. The evidence is consistent with the possibility that RTEL1 promotes replication fork progression *in vivo* by disrupting covalent and/or non-covalent nucleoprotein complexes. However, the production of ssDNA downstream of a DPC by a converging fork or other accessory helicases appears to be sufficient to promote CMG bypass. Therefore, loss of RTEL1 alone may not detectably inhibit DPC repair in cells. Future work will be required to address whether any of the phenotypes observed in RTEL1-deficient mice or humans are attributable to defective bypass of nucleoprotein complexes.

STAR*METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- CONTACT FOR REAGENT AND RESOURCE SHARING
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
 - Xenopus laevis
 - Insect cell lines
- METHODS DETAILS
 - Preparation of DNA constructs
 - *Xenopus* eqq extracts
 - DNA replication in NPE
 - Nascent strand analysis
 - Antibodies and immunodepletion
 - Protein expression and purification
 - Plasmid pull-down
 - DPC pull-down
 - Expressing, and purifying recombinant GINS
 - Purification of Hpall for sortase labeling
 - Fluorescent labeling of GINS and Hpall
 - Preparing substrate with fluorescent DPCs
 - Single-molecule KEHRMIT assay
 - O KEHRMIT data analysis
 - KEHRMIT assay strengths and limitations
- QUANTIFICATION AND STATISTICAL ANALYSIS

SUPPLEMENTAL INFORMATION

Supplemental Information includes seven figures and can be found with this article online at https://doi.org/10.1016/j.cell.2018.10.053.

ACKNOWLEDGMENTS

We thank Niels Mailand, Peter Burgers, Joe Loparo, and the Walter laboratory for feedback on the manuscript, Seungwoo Chang for sortase, Ravi Amunugama for characterizing the FANCM antibody, and Livio Dukaj for Fen1^{mKikGR}. Single-molecule data were collected at the Nikon Imaging Center (NIC) at Harvard Medical School with technical assistance by Jennifer Waters and Talley Lambert. J.L.S. and G.C. were supported by postdoctoral fellowships from the Damon Runyon Cancer Research Foundation and the Jane Coffin Childs Memorial Fund, respectively. J.C.W. is supported by NIH (grants GM80676 and HL098316). J.C.W. is an investigator of the Howard Hughes Medical Institute. J.P.D. received funding from the European Research Council (grant agreement No. 715975) and the Novo Nordisk Foundation (grant agreement NNF14CC0001). M.R. is supported by a grant from the German Research Foundation (Research Group FOR2800).

AUTHOR CONTRIBUTIONS

J.L.S., G.C., and J.C.W. designed and analyzed the experiments and prepared the manuscript. J.L.S. and G.C. performed ensemble and single-molecule experiments, respectively. A.O.G. developed the protocol to methylate DPCs. M.R. and M.M. performed and analyzed the mass spectrometry. J.P.D. and N.B.L. supplied the pDPC^{ssDNA} and pmeDPC^{ssDNA} plasmids. J.P.D. developed the stringent plasmid pull-down protocol.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: March 9, 2018 Revised: September 13, 2018 Accepted: October 24, 2018 Published: December 27, 2018

REFERENCES

Arias, E.E., and Walter, J.C. (2005). Replication-dependent destruction of Cdt1 limits DNA replication to a single round per cell cycle in Xenopus egg extracts. Genes Dev. *19*, 114–126.

Bochman, M.L., and Schwacha, A. (2008). The Mcm2-7 complex has in vitro helicase activity. Mol. Cell *31*, 287–293.

Budzowska, M., Graham, T.G., Sobeck, A., Waga, S., and Watler, J.C. (2015). Regulation of the Rev1-pol ζ complex during bypass of a DNA interstrand cross-link. EMBO J *34*, 1971–1985.

Byun, T.S., Pacek, M., Yee, M.C., Walter, J.C., and Cimprich, K.A. (2005). Functional uncoupling of MCM helicase and DNA polymerase activities activates the ATR-dependent checkpoint. Genes Dev. *19*, 1040–1052.

Chen, I., Dorr, B.M., and Liu, D.R. (2011). A general strategy for the evolution of bond-forming enzymes using yeast display. Proc. Natl. Acad. Sci. USA *108*, 11399–11404.

Dewar, J.M., Budzowska, M., and Walter, J.C. (2015). The mechanism of DNA replication termination in vertebrates. Nature 525, 345–350.

Dewar, J.M., Low, E., Mann, M., Räschle, M., and Walter, J.C. (2017). CRL2^{Lrr1} promotes unloading of the vertebrate replisome from chromatin during replication termination. Genes Dev. *31*, 275–290.

Ding, H., Schertzer, M., Wu, X., Gertsenstein, M., Selig, S., Kammori, M., Pourvali, R., Poon, S., Vulto, I., Chavez, E., et al. (2004). Regulation of murine telomere length by Rtel: an essential gene encoding a helicase-like protein. Cell *117*, 873–886.

Douglas, M.E., Ali, F.A., Costa, A., and Diffley, J.F.X. (2018). The mechanism of eukaryotic CMG helicase activation. Nature 555, 265–268.

Duxin, J.P., Dewar, J.M., Yardimci, H., and Walter, J.C. (2014). Repair of a DNA-protein crosslink by replication-coupled proteolysis. Cell *159*, 346–357.

Duzdevich, D., Warner, M.D., Ticau, S., Ivica, N.A., Bell, S.P., and Greene, E.C. (2015). The dynamics of eukaryotic replication initiation: origin specificity, licensing, and firing at the single-molecule level. Mol. Cell *58*, 483–494.

Fu, Y.V., Yardimci, H., Long, D.T., Ho, T.V., Guainazzi, A., Bermudez, V.P., Hurwitz, J., van Oijen, A., Schärer, O.D., and Walter, J.C. (2011). Selective bypass of a lagging strand roadblock by the eukaryotic replicative DNA helicase. Cell *146*, 931–941.

Fullbright, G., Rycenga, H.B., Gruber, J.D., and Long, D.T. (2016). p97 Promotes a Conserved Mechanism of Helicase Unloading during DNA Cross-Link Repair. Mol. Cell. Biol. *36*, 2983–2994.

Georgescu, R., Yuan, Z., Bai, L., de Luna Almeida Santos, R., Sun, J., Zhang, D., Yurieva, O., Li, H., and O'Donnell, M.E. (2017). Structure of eukaryotic CMG helicase at a replication fork and implications to replisome architecture and origin initiation. Proc. Natl. Acad. Sci. USA *114*, E697–E706.

Graham, J.E., Marians, K.J., and Kowalczykowski, S.C. (2017). Independent and stochastic action of DNA polymerases in the replisome. Cell *169*, 1201–1213.

Guy, C.P., Atkinson, J., Gupta, M.K., Mahdi, A.A., Gwynn, E.J., Rudolph, C.J., Moon, P.B., van Knippenberg, I.C., Cadman, C.J., Dillingham, M.S., et al. (2009). Rep provides a second motor at the replisome to promote duplication of protein-bound DNA. Mol. Cell *36*, 654–666.

Huang, J., Liu, S., Bellani, M.A., Thazhathveetil, A.K., Ling, C., de Winter, J.P., Wang, Y., Wang, W., and Seidman, M.M. (2013). The DNA translocase FANCM/MHF promotes replication traverse of DNA interstrand crosslinks. Mol. Cell *52*, 434–446.

Ide, H., Shoulkamy, M.I., Nakano, T., Miyamoto-Matsubara, M., and Salem, A.M. (2011). Repair and biochemical effects of DNA-protein crosslinks. Mutat. Res. *711*, 113–122.

Ivessa, A.S., Zhou, J.Q., and Zakian, V.A. (2000). The Saccharomyces Pif1p DNA helicase and the highly related Rrm3p have opposite effects on replication fork progression in ribosomal DNA. Cell *100*, 479–489.

Klimasauskas, S., Kumar, S., Roberts, R.J., and Cheng, X. (1994). Hhal methyltransferase flips its target base out of the DNA helix. Cell 76, 357–369.

Langston, L.D., Zhang, D., Yurieva, O., Georgescu, R.E., Finkelstein, J., Yao, N.Y., Indiani, C., and O'Donnell, M.E. (2014). CMG helicase and DNA polymerase ε form a functional 15-subunit holoenzyme for eukaryotic leading-strand DNA replication. Proc. Natl. Acad. Sci. USA *111*, 15390–15395.

Langston, L.D., Mayle, R., Schauer, G.D., Yurieva, O., Zhang, D., Yao, N.Y., Georgescu, R.E., and O'Donnell, M.E. (2017). Mcm10 promotes rapid isomerization of CMG-DNA for replisome bypass of lagging strand DNA blocks. eLife 6, e29118.

Larsen, N.B., Gao, A.O., Sparks, J.L., Gallina, I., Wu, R.A., Mann, M., Räschle, M., Walter, J.C., and Duxin, J.P. (2018). Replication-Coupled DNA-Protein Crosslink Repair by SPRTN and the Proteasome in *Xenopus* Egg Extracts. Mol Cell 73. https://doi.org/10.1016/j.molcel.2018.11.024.

Lebofsky, R., van Oijen, A.M., and Walter, J.C. (2011). DNA is a co-factor for its own replication in Xenopus egg extracts. Nucleic Acids Res. 39, 545–555.

Lessel, D., Vaz, B., Halder, S., Lockhart, P.J., Marinovic-Terzic, I., Lopez-Mosqueda, J., Philipp, M., Sim, J.C., Smith, K.R., Oehler, J., et al. (2014). Mutations in SPRTN cause early onset hepatocellular carcinoma, genomic instability and progeroid features. Nat. Genet. *46*, 1239–1244.

Lewis, J.S., Spenkelink, L.M., Schauer, G.D., Hill, F.R., Georgescu, R.E., O'Donnell, M.E., and van Oijen, A.M. (2017). Single-molecule visualization of *Saccharomyces cerevisiae* leading-strand synthesis reveals dynamic interaction between MTC and the replisome. Proc. Natl. Acad. Sci. USA *114*, 10630–10635.

Ling, C., Huang, J., Yan, Z., Li, Y., Ohzeki, M., Ishiai, M., Xu, D., Takata, M., Seidman, M., and Wang, W. (2016). Bloom syndrome complex promotes FANCM recruitment to stalled replication forks and facilitates both repair and traverse of DNA interstrand crosslinks. Cell Discov. *2*, 16047.

Lopez-Mosqueda, J., Maddi, K., Prgomet, S., Kalayil, S., Marinovic-Terzic, I., Terzic, J., and Dikic, I. (2016). SPRTN is a mammalian DNA-binding metalloprotease that resolves DNA-protein crosslinks. eLife 5, e21491.

Loveland, A.B., Habuchi, S., Walter, J.C., and van Oijen, A.M. (2012). A general approach to break the concentration barrier in single-molecule imaging. Nat. Methods *9*, 987–992.

Maric, M., Maculins, T., De Piccoli, G., and Labib, K. (2014). Cdc48 and a ubiquitin ligase drive disassembly of the CMG helicase at the end of DNA replication. Science *346*, 1253596. Maskey, R.S., Kim, M.S., Baker, D.J., Childs, B., Malureanu, L.A., Jeganathan, K.B., Machida, Y., van Deursen, J.M., and Machida, Y.J. (2014). Spartan deficiency causes genomic instability and progeroid phenotypes. Nat. Commun. 5, 5744.

Moreno, S.P., Bailey, R., Campion, N., Herron, S., and Gambus, A. (2014). Polyubiquitylation drives replisome disassembly at the termination of DNA replication. Science *346*, 477–481.

O'Donnell, M.E., and Li, H. (2018). The ring-shaped hexameric helicases that function at DNA replication forks. Nat. Struct. Mol. Biol. 25, 122–130.

Samel, S.A., Fernández-Cid, A., Sun, J., Riera, A., Tognetti, S., Herrera, M.C., Li, H., and Speck, C. (2014). A unique DNA entry gate serves for regulated loading of the eukaryotic replicative helicase MCM2-7 onto DNA. Genes Dev. 28, 1653–1666.

Sparks, J., and Walter, J.C. (2018). Extracts for analysis of DNA replication in a nucleus-free system. Cold Spring Harb. Protoc. Published online May 16, 2018. https://doi.org/10.1101/pdb.prot097154.

Stano, N.M., Jeong, Y.J., Donmez, I., Tummalapalli, P., Levin, M.K., and Patel, S.S. (2005). DNA synthesis provides the driving force to accelerate DNA unwinding by a helicase. Nature *435*, 370–373.

Stingele, J., Schwarz, M.S., Bloemeke, N., Wolf, P.G., and Jentsch, S. (2014). A DNA-dependent protease involved in DNA-protein crosslink repair. Cell *158*, 327–338.

Stingele, J., Bellelli, R., Alte, F., Hewitt, G., Sarek, G., Maslen, S.L., Tsutakawa, S.E., Borg, A., Kjær, S., Tainer, J.A., et al. (2016). Mechanism and regulation of DNA-protein crosslink repair by the DNA-dependent metalloprotease SPRTN. Mol. Cell *64*, 688–703.

Stingele, J., Bellelli, R., and Boulton, S.J. (2017). Mechanisms of DNA-protein crosslink repair. Nat. Rev. Mol. Cell Biol. *18*, 563–573.

Taylor, M.R.G., and Yeeles, J.T.P. (2018). The initial response of a eukaryotic replisome to DNA damage. Mol. Cell *70*, 1067–1080.

Trowitzsch, S., Bieniossek, C., Nie, Y., Garzoni, F., and Berger, I. (2010). New baculovirus expression tools for recombinant protein complex production. J. Struct. Biol. *172*, 45–54.

Vannier, J.B., Sandhu, S., Petalcorin, M.I., Wu, X., Nabi, Z., Ding, H., and Boulton, S.J. (2013). RTEL1 is a replisome-associated helicase that promotes telomere and genome-wide replication. Science *342*, 239–242.

Vannier, J.B., Sarek, G., and Boulton, S.J. (2014). RTEL1: functions of a disease-associated helicase. Trends Cell Biol. 24, 416–425.

Vaz, B., Popovic, M., Newman, J.A., Fielden, J., Aitkenhead, H., Halder, S., Singh, A.N., Vendrell, I., Fischer, R., Torrecilla, I., et al. (2016). Metalloprotease SPRTN/DVC1 orchestrates replication-coupled DNA-protein crosslink repair. Mol. Cell *64*, 704–719.

Walter, J., and Newport, J.W. (1997). Regulation of replicon size in Xenopus egg extracts. Science 275, 993–995.

Walter, J., and Newport, J. (2000). Initiation of eukaryotic DNA replication: origin unwinding and sequential chromatin association of Cdc45, RPA, and DNA polymerase alpha. Mol. Cell *5*, 617–627.

Yardimci, H., Loveland, A.B., van Oijen, A.M., and Walter, J.C. (2012a). Singlemolecule analysis of DNA replication in Xenopus egg extracts. Methods 57, 179–186.

Yardimci, H., Wang, X., Loveland, A.B., Tappin, I., Rudner, D.Z., Hurwitz, J., van Oijen, A.M., and Walter, J.C. (2012b). Bypass of a protein barrier by a replicative DNA helicase. Nature *492*, 205–209.

STAR*METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit Anti-xIRTEL1-N; Antigen: amino acids 400-654	This study	N/A
Rabbit Anti-xIRTEL1-C; Antigen: HPDTSQRKKPRGDIWSC	This study	N/A
Rabbit Anti-FancM; Antigen: CYMFEEEMVPPENPTKRSRVS	This study	N/A
Rabbit Anti-MCM6; Antigen: CLVVNPNYMLED	This study	N/A
Rabbit Anti-GINS; Antigen: purified GINS complex	This study	N/A
Rabbit Anti-Orc2	Walter and Newport, 1997	N/A
Rabbit Anti-CDC45	Walter and Newport, 2000	N/A
Rabbit Anti-M.Hpall	Larsen et al., 2018	N/A
Rabbit Anti-PSMA3	Larsen et al., 2018	N/A
Rabbit Anti-SPRTN-N	Larsen et al., 2018	N/A
Rabbit Anti-Chk1-p(Ser345)	Cell Signaling	Cat# 2341L; RRID: AB_330023
Rabbit Anti-Histone H3	Cell Signaling	Cat# 9715S; RRID: AB_331563
Rabbit serum	Sigma	Cat# R9133-5ML
Bacterial and Virus Strains	,	
T7 Express	NEB	Cat# C2566I
DH10β	NEB	Cat# C3019I
DH10EMBacY	Geneva Biotech	Cat# DH10EMBacY
Chemicals, Peptides, and Recombinant Proteins		
[α- ³² P] dATP	Perkin Elmer	Cat# BLU512H500UC
biotin-dCTP	Thermo Fisher	Cat# 19518018
biotin-dGTP	Perkin Elmer	Cat# NEL541001EA
SAM	NEB	Cat# B9003S
ATP	Sigma	Cat# A-5394
Phosphocreatine	Sigma	Cat# P-6502
Creatine Phosphokinase	Sigma	Cat# C-3755
Nucleoplasmic Extract (NPE)	Sparks and Walter, 2018	N/A
High Speed Supernatant (HSS)	Sparks and Walter, 2018	N/A
Geminin	Arias and Walter, 2005	N/A
ProteinaseK	Roche	Cat# 3115879001
RNaseA	Sigma	Cat# R4642-250mg
HRV 3C protease (Thermo Fisher)	Thermo Fisher	Cat# 88946
GINS expressed from pGC128	This study	N/A
Fen1 ^{mKikGR}	Loveland et al., 2012	N/A
M.Hpall-His ₆	Duxin et al., 2014	N/A
M.Hpall-10aa-LPETG-His ₆ expressed from pGC220	This study	N/A
Lacl-biotin	Duxin et al., 2014	N/A
Lacl-His ₆	Duxin et al., 2014	N/A
Sortase	Chen et al., 2011, Gift from Seungwoo Chang	N/A
p97i (NMS873)	Sigma	Cat# SML1128
MG262	Boston Biochem	Cat# I-120
Cdc7-i (PHA-767491)	Sigma	Cat# PZ0178
ATRi (ETP-46464)	Sigma	Cat# SML1321

(Continued on next page)

Continued		
REAGENT or RESOURCE	SOURCE	IDENTIFIER
Aphidicolin	Sigma	Cat# A0781
IPTG	Sigma	Cat# 15502
Leupeptin	Roche	Cat# 11529048001
Aprotinin	Roche	Cat# 11583794001
Triton X-100	Sigma	Cat# X100-100ML
BSA	Fisher	Cat# BP1600-100
Hoechst solution	Fisher	Cat# 62249
L cysteine	Fisher	Cat# ICN10144601
Cycloheximide	Sigma	Cat# C7698-5G
Cytochalasin B	Sigma	Cat# C6762-10MG
14-ml Round-Bottom Falcon Tubes	Fisher	Cat# 352059
2.5-ml Thin-Walled Ultracentrifuge Tubes	Beckman	Cat# 347356
Nocodazole	Sigma	Cat# M1404-10MG
EDTA-free cOmplete protease inhibitor cocktail	Roche	Cat# 11873580001
Nb.Bsml	NEB	Cat# R0706S
Notl-HF	NEB	Cat# R3189S
Aatll	NEB	Cat# R0117S
Fspl	NEB	Cat# R0135S
Nt.Bbvcl	NEB	Cat# R0632S
Nb.Bbvcl	NEB	Cat# R0631S
Nt.BspQI	NEB	Cat# R0644S
Xmnl	NEB	Cat# R0194S
Klenow Fragment $3' \rightarrow 5'$ exo-	NEB	Cat# M0212S
AlexaFluor647-maleimide	Thermo Fisher	Cat# A20347
AlexaFluor568-maleimide	Thermo Fisher	Cat# 20341
SYBR Safe	Thermo Fisher	Cat# S33102
SYTOX Green	Thermo Fisher	Cat# S7020
GGGGYKCK peptide	New England Peptide	N/A
HPDTSQRKKPRGDIWSC peptide	New England Peptide	N/A
CYMFEEEMVPPENPTKRSRVS peptide	Bethyl Laboratories	N/A
PD10 desalting column	GE Healthcare	Cat# 17-0851-01
NiNTA resin	QIAGEN	Cat# 30410
Streptavidin-coupled magnetic Dynabeads M-280	Invitrogen	Cat# 11206D
Glutatione Sepharose 4B	GE Healthcare	Cat# 17075605
Protein A Sepharose Fast Flow	GE Healthcare	Cat# 17127903
AminoLink Coupling Resin	Thermo Fisher	Cat# 20381
Double-sided tape	Grace Bio-Labs	Cat# 620001
25x75x1mm slides	VWR	Cat# 48300-025
24x60mm No. 1.5 coverslips	VWB	Cat# 16004-312
Biotin-PEG-SVA	Lavsan Bio	Cat# Bio-PEG-SVA-5k-100mg
M-PEG-SVA MW5000	Lavsan Bio	Cat# MPEG-SVA-5K-1g
Critical Commercial Assays		
Bac-to-Bac Baculovirus Expression System	Thermo Fisher	Cat# 10359016
MultiBac Expression System Kit	Geneva Biotech	Cat# MultiBac
ZB BAC DNA minipren kit	Zymo Research	Cat# D4048
Evperimental Models: Cell Lines		
	Expression Systems	Cat# 94-001S
	Expression Systems	Cat# 94-0015
Thi insect cells	Expression Systems	Gal# 94-0025

(Continued on next page)

Continued		
REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains		
Xenopus laevis, adult female	Nasco	Cat# LM00535
Xenopus laevis, adult male	Nasco	Cat# LM00715
Oligonucleotides		
5'GGGAGCTGAATGCCGCGCGAATAATGGTTTCTTAGACGT3'	IDT	N/A
5'CATCCACTAGCCAATTTATGCTGAGGTACCGGATTGAGTAGC	IDT	N/A
TACCGGATGCTGAGGGGATCCACTAGCCAATTTATCATGG3'		
5'cgcgcttaatcagtgaggcacctatctcCGGTCTGAGTCATGCGTAActc gagtGCTTGTAGTGGATTtaCCGGATtgagtagcTACCGGATggtac3'	IDT	N/A
5'CATCCGGTAGCTACTCAATCCGGTAAATCCACTACAAGCACT CGAGTTACGCATGACTCAGACCGGAGATAGGTGCCTCACTGA TTAAG3'	IDT	N/A
5'cATCCGGTAgctactcaATCCGGtaTTAGGTGATGTTCGtgagctc AATGCGTACTGAGTCTGGCgagataggtgcctcactgattaag3'	IDT	N/A
5'TCAGCATCCGGTAGCTACTCAATC[C5FluordC]GGTACC3'	Bio-Synthesis	N/A
5'TCAGCATC[C5FluordC]GGTAGCTACTCAATCCGGTACC3'	Bio-Synthesis	N/A
5'tgaggTAC[C5FluordC]GGATtgagtagcTACCGGATgc-3'	Bio-Synthesis	N/A
5'TCAGCATCCGGTAGCTACTCAATC[C5FluordC]GGTACC3'	Bio-Synthesis	N/A
5'tgaggTACCGGATtgagtagcTAC[C5FluordC]GGATgc3'	Bio-Synthesis	N/A
5'TCAGCATC[C5FluordC]GGTAGCTACTCAATC[C5FluordC]GG TACC3'	Bio-Synthesis	N/A
Recombinant DNA	,	
pDPC ^{Lead} (same as pJLS2)	This study	N/A
pDPC ^{Lag} (same as pJLS2)	This study	N/A
pJD2	Duxin et al., 2014	N/A
pDPC ^{2xLead}	This study	N/A
pDPC ^{Lag/Lead}	This study	N/A
pDPC ^{Lag/Lead-Bubble}	This study	N/A
pJLS3	This study	N/A
pDPC ^{ssDNA}	Larsen et al., 2018	N/A
pLacO ₃₂	Dewar et al., 2015	N/A
pJLS100	This study	N/A
pJLS101	This study	N/A
pGC128	This study	N/A
pGC220	This study	N/A
pGC261	This study	N/A
λ DNA	NEB	Cat# N3011S
pBlueScript	Agilent	Cat# 212205
Software and Algorithms		
Multi-Gauge	Fujifilm	N/A
NIS Elements	Nikon Instruments	https://www.nikoninstruments.com/ Products/Software
MATLAB	Mathworks	https://www.mathworks.com/ products/matlab.html
ImageJ	NIH	https://imagej.nih.gov/ij/download.html
Image Stabilizer Plugin for ImageJ	Kang Li	http://www.cs.cmu.edu/~kangli/ code/Image_Stabilizer.html

CONTACT FOR REAGENT AND RESOURCE SHARING

Requests for resources, reagents, and further information about the methods should be directed to the lead contact, Johannes Walter (johannes_walter@hms.harvard.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Xenopus laevis

Egg extracts were prepared using eggs from adult *Xenopus laevis* female frogs (Nasco Cat# LM00535) and sperm chromatin was prepared from the testes of adult *Xenopus laevis* male frogs (Nasco Cat #LM00715). All animals were healthy and not previously subjected to any other procedures or immunizations. No animal husbandry was performed for this study. Frogs were housed at the Satellite Amphibian Facility of the BCMP Department at Harvard Medical School in compliance with Institutional Animal Care and Use Committee (IACUC) regulations. All experiments involving animals were approved by the Harvard Medical Area IACUC and conform to the relevant regulatory standards.

Insect cell lines

Sf9 cells (Expression Systems Cat# 94-001S) and Tni cells (Expression Systems Cat# 94-002S) were cultured at 27°C for protein overexpression. Cells were cultured in ESF 921 insect cell culture medium (Fisher scientific Cat#96-001-01-CS).

METHODS DETAILS

Preparation of DNA constructs

To generate pDPC, we first created pJLS2 by replacing the AatII-BsmBI fragment from pJD2 (Duxin et al., 2014) with the a sequence (5'-GGGAGCTGAATGCCGCGCGAATAATGGTTTCTTAGACGT-3') which contains a Nb.Bsml site. To generate pDPC^{2xLead} the Sacl-BssHII fragment from pJLS2 was replaced with the following sequence: 5'- CATCCACTAGCCAATTTATGCTGAGGTACCGGATT GAGTAGCTACCGGATGCTGAGGGGATCCACTAGCCAATTTATCATGG-3'. To generate pDPC^{Lag/Lead}, and pDPC^{Lag/Lead-Bubble} we first replaced the BssHII-KpnI fragment from pJLS2 with the sequence 5'- CGCGCTTAATCAGTGAGGCACCTATCTCCGGTCT GAGTCATGCGTAACTCGAGTGCTTGTAGTGGATTTAC[C5-Fluor dC]GGATTGAGTAGCTACCGGATGGTAC-3' hybridized with either 5'- CATC[C5-Fluor dC] GGTAGCTACTCAATCCGGTAAATCCACTACAAGCACTCGAGTTACGCATGACTCAGACCGGAGA TAGGTGCCTCACTGATTAAG-3' or 5'-CATC/C5-Fluor dC/GGTAGCTACTCAATCCGGTATTAGGTGATGTTCGTGAGCTCAATGCG TACTGAGTCTGGCGAGATAGGTGCCTCACTGATTAAG -3', respectively. The supercoiled band was purified by cesium chloride gradient ultracentrifugation. The C5-Fluor dC modified plasmids were mixed with either methylated M.Hpall or nonmethylated M.Hpall in M.Hpall reaction buffer (50 mM Tris-HCl, pH 7.5, 5 mM 2-mercaptoethanol, 10 mM EDTA) and supplemented with 100 mM S-adenosylmethionine (NEB, Ipswich, MA) for 12-18 hr at 37°C. pJLS2 or pJLS3 were nicked with Nt.Bbvcl and ligated with an oligonucleotide containing a fluorinated cytosine (5'-TCAGCATCCGGTAGCTACTCAATC[C5-Fluor dC]GGTACC-3') and subsequently crosslinked to M.Hpall-His₆ or methylated M.Hpall-His₆ to generate pDPC^{Lead} and pDPC^{2xLead} or pmeDPC^{Lead} and pmeDPC^{2xLead}, respectively, as previously described (Duxin et al., 2014). pJLS2 was nicked with Nb.Bbvcl and ligated with an oligonucleotide containing a fluorinated cytosine (5'-TCAGCATC[C5-Fluor dC]GGTAGCTACTCAATCCGGTACC-3') and subsequently crosslinked to methylated M.Hpall-His₆ to generate pmeDPC^{Lag}, as previously described (Duxin et al., 2014). To create pmeDPC^{Lead/Lag}, pJLS2 was first nicked with Nt.Bbvcl and ligated with an oligonucleotide containing a fluorinated cytosine (5'-TCAGCATC[C5-Fluor dC]GGTAGCTACTCAATCCGGTACC-3'). It was subsequently nicked with Nb.Bbvcl and ligated with a second oligonucleotide containing a fluorinated cytosine (5'- TGAGGTAC[C5-Fluor dC]GGATTGAGTAGCTACCGGATGC-3') before crosslinking to methylated M.Hpall-His₆. To create pmeDPC^{Lag/Lead} pJLS2 was first nicked with Nt.Bbvcl and ligated with an oligonucleotide containing a fluorinated cytosine (5'-TCAGCATCCGGTAGCTACTCAATC[C5-Fluor dC]GGTACC-3'). It was subsequently nicked with Nb.Bbvcl and ligated with a second oligonucleotide containing a fluorinated cytosine (5'- TGAGGTACCGGATTGAGTAG CTAC[C5-Fluor dC]GGATGC-3') before crosslinking to methylated M.Hpall-His₆. To create pmeDPC^{Lead/Lead} or pmeDPC^{2xLeadLead}, pJLS2 or pJLS3 was nicked with Nt.Bbvcl and ligated with an oligonucleotide containing two fluorinated cytosines (5'-TCAGCATC [C5-Fluro dC]GGTAGCTACTCAATC[C5-Fluro dC]GGTACC-3') and subsequently crosslinked to methylated M.Hpall-His₆ to generate pmeDPC^{Lead/Lead} or pDPC^{2xLeadLead}, respectively, as previously described (Duxin et al., 2014). Creation of pDPC^{ssDNA} and pmeDPC^{ssDNA} is described in Larsen et al. (2018). Creation of pLacO₃₂ was previously described (Dewar et al., 2015).

Xenopus egg extracts

equation reference goes hereFor detailed protocols for preparation of demembranated sperm chromatin, High-Speed Supernatant (HSS), and Nucleoplasmic Extract (NPE) see Sparks and Walter (2018).

Preparation of demembranate sperm chromatin

Testes are harvested from 6-8 adult male *Xenopus laevis* frogs, and minced into tiny pieces using a clean razor blade while immersed in 1mL buffer #1 (10 mM HEPES pH 7.4, 80 mM KCl, 15 mM NaCl, 5 mM MgCl₂, 1 mM EDTA, 0.2 M sucrose). The sperm are released by vortexing. The tissue is pelleted by centrifugation, and the supernatant is collected. Sperm are pelleted by centrifugation at 2600*g*

for 15 minutes at 4°C in a Sorvall Lynx 4000 centrifuge with swinging-bucket rotor TH13-6X50 (or equivalent). In all subsequent steps, the sperm should be kept on ice unless otherwise stated. The supernatant is discarded and the sperm pellet is re-suspended in buffer #4 (10 mM HEPES pH 7.4, 80 mM KCI, 15 mM NaCI, 5 mM MgCl₂, 1 mM EDTA, 2.0 M sucrose). The sperm is isolated on a sucrose step gradient in 2.5 mL thin-walled ultracentrifuge tubes (Beckman Cat# 347356): the upper sucrose layer consists of 1.7 mL Buffer #2 (10 mM HEPES pH 7.4, 80 mM KCI, 15 mM NaCI, 5 mM MgCl₂, 1 mM EDTA, 2.3 M sucrose), and the bottom sucrose layer consists of 0.25 mL Buffer #3 (10 mM HEPES pH 7.4, 80 mM KCl, 15 mM NaCl, 5 mM MgCl₂, 1 mM EDTA, 2.5 M sucrose). The sperm is layered on top of the sucrose cushion and spun down for 45 minutes at 93,000g at 2°C in a tabletop ultracentrifuge (Beckman Optima Max-E or equivalent) in a swing-bucket rotor (Beckman TLS-55 or equivalent). Following centrifugation, the sperm forms a pellet at the bottom of the tube. The top half of the gradient is gently aspirated and discarded, whereas the lower half of the gradient is harvested in a clean tube and combined with the sperm pellet re-suspended in buffer #1. The sperm is diluted in buffer #1 to 12 mL and pelleted by centrifugation in a swing-bucket rotor at 3000g at 4°C for 15 minutes. The supernatant is gently removed and the sperm is re-suspended in buffer #1 supplemented with 1mM DTT, 10 µg/ml of leupeptin (Roche Cat# 11529048001), and 10 µg/ml of aprotinin (Roche Cat# 11583794001). The sperm chromatin is de-membranated by addition of 0.4% Triton X-100 (Sigma Cat# X100-100ML) and incubation with rotation at 4°C for 45 minutes. Detergent is removed by pelleting the sperm through a 0.5 mL cushion of buffer #5 (10 mM HEPES pH 7.4, 80 mM KCl, 15 mM NaCl, 5 mM MgCl₂, 1 mM EDTA, 0.5 M sucrose, 3.0% BSA (Fisher Cat# BP1600-100), 10 µg/ml aprotinin, 10 µg/ml leupeptin, 1 mM DTT). The de-membranated sperm is layered on top of the cushion and spun down for 10 minutes at 750g at room temperature in a benchtop centrifuge with swing-bucket rotor. The supernatant is discarded and the sperm is re-suspended in 0.5ml of buffer #6 (10 mM HEPES pH 7.4, 80 mM KCl, 15 mM NaCl, 5 mM MgCl₂, 1 mM EDTA, 0.2 M sucrose, 3.0% BSA, 10 µg/ml aprotinin, 10 µg/ml leupeptin, 1 mM DTT). The sperm is pelleted for 10 minutes at 750q, and the wash procedure is repeated with buffer #6. Finally, the sperm is resuspended in 1.0ml of buffer #6 and sperm concentration is determined by adding Hoechst solution (Fisher Cat# 62249) and counting the number of sperm on a hemo-cytometer using the UV/DAPI filters on an epi-fluorescence microscope. The preparation is diluted to a final concentration of 220,000sperm/µl and 90µl aliquots are snap frozen in liquid nitrogen for subsequent use in preparing Nucleoplasmic Extract (NPE).

Preparation of High-Speed Supernatant

Ovulation is induced in six adult female *Xenopus laevis* frogs according to standard protocols. *Xenopus* eggs are laid in water supplemented with 100 mM NaCl and harvested after 20-22 hours. Eggs are de-jellied in 1 l of 2.2% L cysteine pH 7.7 (Fisher Cat# ICN10144601), washed with 2 l of 0.5 X Marc's Modified Ringer's (MMR) (2.5 mM HEPES pH 7.8, 50 mM NaCl, 1 mM KCl, 0.25 mM MgSO₄, 1.25 mM CaCl₂, 0.05 mM EDTA), then washed with 1 l of Egg Lysis Buffer (ELB) sucrose (10 mM HEPES pH 7.7, 50 mM KCl, 2.5 mM MgCl₂, 250 mM sucrose, 1 mM DTT, and 50 μ g/ml cycloheximide (Sigma Cat# C7698-5G). Eggs are packed in 14 mL round-bottom Falcon tubes (Fisher Cat# 352059) by centrifugation for 1 minute at 200g in a swing-bucket benchtop centrifuge and excess buffer is aspirated. The packed eggs are supplemented with 5 μ g/ml aprotinin, 5 μ g/ml leupeptin, and 2.5 μ g/ml cyclohalasin B (Sigma Cat# C6762-10MG) and crushed by centrifugation for 20 minutes at 20,000g at 4°C in a swinging-bucket rotor in a Sorvall Lynx 4000 centrifuge (or equivalent). It is essential that the rotors be kept at room temperature until the centrifugation. After centrifugation the crushed eggs are kept on ice. The top layer of low-speed supernatant (LSS) is collected and supplemented with 50 μ g/ml cycloheximide, 1 mM DTT, 10 μ g/ml aprotinin, 10 μ g/ml leupeptin, and 5 μ g/ml cytochalasin B. The LSS is transferred to 2.5 mL thin-walled ultracentrifuge tubes (Beckman Cat# 347356) and centrifuged for 90 minutes at 260,000g at 2°C in a tabletop ultracentrifuge with a swing-bucket rotor. After centrifugation the clear high-speed supernatant (HSS) layer is harvested, snap frozen in liquid nitrogen, and stored at -80°C.

Preparation of Nucleoplasmic Extract

Eggs from 20 adult female frogs are collected, washed, and de-jellied as described above for HSS, except use double the volume of L-cysteine, 0.5xMMR, and ELB-sucrose. Prepare LSS as described above, and supplement LSS with 50 μ g/ml cycloheximide, 1 mM DTT, 10 μ g/ml aprotinin, 10 μ g/ml leupeptin, 5 μ g/ml cytochalasin B, and 3.3 μ g/ml nocodazole (Sigma Cat# M1404-10MG). The extract is spun down for 15 minutes at 20,000g in a Sorvall Lynx 4000 centrifuge with swing-bucket rotor. After centrifugation the top lipid layer is aspirated and the extract is decanted into a clean tube leaving behind the black pellet. The extract is warmed up to room temperature and supplemented with ATP regeneration system: 2 mM ATP, 20 mM phosphocreatine, and 5 μ g/ml phosphokinase. Nuclei are assembled by thoroughly mixing 4.5ml of extract with one 90 μ l aliquot of demembranated sperm chromatin and incubating at room temperature for 75-90 minutes. The nuclear assembly reaction is centrifuged for 2.5 minutes at 20,000g at 4°C in a Sorvall Lynx 4000 in a swing-bucket rotor. The layer of nuclei formed at the top is collected and transferred to a clean tube and clarified by centrifugation at 260,000g at 2°C for 30 minutes. Following centrifugation, the top layer of lipids is carefully aspirated and the clear nucleoplasmic extract (NPE) is harvested carefully to avoid contamination with the chromatin pellet. The NPE is snap frozen in liquid nitrogen and stored at -80° C.

DNA replication in NPE

Plasmid DNA replication in *Xenopus* egg extract is carried out by supplementing high-speed supernatant (HSS) of egg cytoplasm with plasmid DNA at a final concentration of 7.5–15 ng/ μ L (Licensing reaction). Licensing reactions were incubated for 30 min at room temperature to assemble pre-replicative complexes (pre-RCs). To prevent licensing, Geminin was added to HSS at a final concentration of 10 μ M and incubated for 10 min at room temperature prior to addition of plasmid DNA. To initiate DNA replication, 1 volume of licensing reaction was mixed with 2 volumes of nucleoplasmic extract (NPE) that had been diluted two-fold with

1xELB-sucrose (10 mM HEPES-KOH pH 7.7, 2.5 mM MgCl₂, 50 mM KCl, 250 mM sucrose). For radiolabeling DNA replication products, $[\alpha$ -³²P] dATP was added to HSS prior to the addition of plasmid DNA. For replication in the presence of Lacl, 1 volume of plasmid (75 ng/µL) was incubated with an equal volume of 12 µM LacI for 30 minutes prior to transfer into HSS so that the final concentration of plasmid was 7.5 ng/µl (Duxin et al., 2014). For native agarose gel electrophoresis 0.5 µl aliquots of replication reaction were typically stopped with 5–10 volumes of replication stop buffer (8 mM EDTA, 0.13% phosphoric acid, 10% ficoll, 5% SDS, 0.2% bromophenol blue, 80 mM Tris-HCl at pH 8), treated with 1 µg/µL Proteinase K for 1 hr at 37°C, and applied to a 0.8 agararose gel. For nascent strand analysis, 2.5 µl aliquots of replication reaction were stopped in 10 volumes of sequencing stop buffer (0.5% SDS, 25 mM EDTA, 50 mM Tris-HCl pH 8.0) followed by addition of 1.25 µl of 190 ng/µL RNase A and incubated for 30 minutes at 37°C. After RNase digestion, 1.25 µl of 900 ng/µL Proteinase K was added to the DNA samples and incubated overnight at room temperature. Following the Proteinase K treatment, samples were diluted to 150 µl with 10 mM Tris-HCl pH 8.0. The samples were extracted once with an equal volume of phenol/chloroform followed by one extraction with an equal volume of chloroform. The DNA was then precipitated with the addition of 0.1 volumes 3M sodium acetate pH 5.2 and 1 µl glycogen (20 mg/ml stock) and resuspended in 7.5 µl. For RTEL1 immunodepletion and rescue experiments, NPE was supplemented with ~200 nM recombinant wild-type or mutant Xenopus RTEL1 and incubated for 15 minutes prior to replication initiation. For p97i (NMS873; Sigma Cat# SML1128) treatment, NPE was supplemented with 200 µM NMS-873 (20 mM stock) and incubated for 10 minutes prior to mixing with HSS (133.33 µM final concentration in replication mix. For MG262 (stock 20 mM; Boston Biochem.Cat# I-120) treatment, NPE was supplement with 200 μM MG262 and incubated for 15 minutes prior to mixing with HSS (133.33 µM final concentration in replication mix). A 1 mM Cdc7-i (PHA-767491; Sigma Cat# PZ0178) stock was prepared in ELB-sucrose buffer and added to replication mix at a final working concentration of 100 µM at the specified time point. For ATRi (ETP-46464; Sigma Cat# SML1321) treatment, NPE was supplemented with 100 µM ATRi and incubated for 10 minutes prior to mixing with HSS (66.67 µM final concentration in replication mix). For aphidicolin (Sigma Cat# A0781) treatment, replication mix was supplemented with 150 μM aphidicolin. For IPTG (stock 1 M; Sigma Cat# I5502) treatment, replication reactions were supplemented with 300 µM IPTG 5 minutes after replication initiation. Samples were analyzed by native 0.8% agarose gel electrophoresis. Gels were exposed to phosphorscreens and imaged on a Typhoon FLA 7000 phosphorimager (GE Healthcare). To better visualize products in Figure 3D (bottom panel), the original images were converted into a log scale for display by applying the function $f(p) = log(p)^{*}255/log (255)$ to each pixel (p) in the images.

Nascent strand analysis

To nick radio-labeled nascent leading-strands, 3-4 μ l of extracted and ethanol precipitated DNA (see above) at 1-2 ng μ l⁻¹ was incubated in 1x buffer 3.1 (New England BioLabs) with 0.45 units μ l⁻¹ Nb.Bsml (New England BioLabs) in a 5 μ l reaction at 65°C for 1 h. To digest radio-labeled nascent leading-strand 3-4 μ l of extracted and ethanol precipitated DNA a 1-2 ng μ l⁻¹ was incubated in 1x cutsmart buffer (New England BioLabs) with 1 unit μ l⁻¹ Aatll (New England BioLabs) and Fspl (New England BioLabs) in a 5 μ l reaction at 37°C for 2 h. To nick rightward leading strands of pLacO₃₂, 3-4 μ l of purified DNA at 1-2 ng μ l⁻¹ was incubated in buffer 3.1 with 0.4 units μ l⁻¹ Nt.BspQI (New England BioLabs) at 37°C for 1 h. Digestion reactions were stopped with 0.5 volumes of Sequencing Stop solution (95% formamide, 20 mM EDTA, 0.05% bromophenol blue, 0.05% xylene cyanol FF). Nicked DNA (3.5 to 4 μ l samples) was separated on 4% (for pLacO) or 7% (pDPC) polyacrylamide sequencing gels. Gels were dried and subjected to phosphorimaging using a Typhoon FLA 7000 phosphoimager. Gels were quantified using Multi Gauge software (Fuji Photo Film). For Figure 6A and 1 μ l of purified DNA was used for Xmnl digestion.

After leading strands stall at the -30 to -44 positions, they approach to the -1 to +1 positions relative to the DPC. Intermediates between these two clusters are also observed, but not when the DPC cannot be ubiquitylated (e.g., Figure 1C) (Duxin et al., 2014). The question arose whether these intermediates are associated with CMGs that have bypassed or not. Importantly, our single molecule data demonstrates that CMG bypasses a DPC with similar kinetics whether or not the DPC is ubiquitylated (Figure S6J). These results indicate that these approach intermediates are generated after CMG bypass, probaly due to polymerase stalling at ubiquitylated DPCs. Therefore, to quantify the percentage of CMG that underwent bypass in Figures 1C, 2, 3C, 3E, 4B, S3D, and S3E (called "approach" in Figure 1C, where bypass had not yet been established), the radioactive signal of all leading strands located between positions +1 and -29 on the gel (reflecting CMGs that have bypassed) was divided by the radioactive signal for leading strands between positions +1 and -44 (reflecting CMGs that have stalled at the lesion or undergone bypass). In the case of pmeDPC^{Lead/Lead} (Figure 2, lanes 13-18), we divided the signal between +1 of the 2nd DPC and -1 of the 1st DPC (both DPCs bypassed) by the signal between +1 of the 2nd DPC and -44 of the first DPC (bypassed and not bypassed).

Antibodies and immunodepletion

The xIRTEL1-N antibody was raised against a fragment of *Xenopus laevis* RTEL1 encompassing amino acids 400-654, which was tagged on its N terminus with His₆ (pJLS100). The protein fragment was overexpressed and purified from bacteria under denaturing conditions, and the antibody was raised by Pocono Rabbit Farm & Laboratory. The RTEL1 antibody was affinity purified from the serum using the RTEL1 antigen according to standard protocols. The xIRTEL1-C antibody was raised against amino acids 428-443 (Ac-HPDTSQRKKPRGDIWSC-amide) by New England Peptide. The FancM antibody was raised against (Ac-CYMFEEEMVPPENPTKRSRVS-amide) FancM peptide by Bethyl Laboratories. The following antibodies were described previously: Orc2 (Walter and Newport, 1997), CDC45 (Walter and Newport, 2000), M.Hpall (Larsen et al., 2018), PSMA3 (Larsen et al., 2018), SPRTN-N (Larsen et al., 2018), Chk1-p(Ser345) (Cell Signaling Cat #2341L), and Histone H3 (Cell Signaling Cat #9715S). Mcm6

antibody was raised against a C-terminal peptide (Ac-CLVVNPNYMLED-OH) and affinity purified. The most prominent band it recognized in western blotting of total extract was 115 kD, and it recognized the same band in protein preparations containing recombinant MCM6. Two rabbits (#34299 and #34300) were immunized with purified GINS (Pocono Rabbit Farm and Laboratory). Anti-GINS antibodies were affinity-purified from serum using GINS immobilized on AminoLink Coupling Resin (Thermo Fisher Cat# 20381). 2-4mg of purified GINS was cross-linked to 1mL of resin according to the manufacturer's protocol. Purified antibody was dialyzed into 1x TBS buffer, concentrated to 1mg/mL final concentration, and stored at -80C.

For RTEL1 immunodepletion, 3.5 volumes of purified RTEL1 antibody (1 mg mL⁻¹) or an equivalent amount of rabbit IgG purified from non-immunized rabbit serum (Sigma) were incubated with 1 volume of Protein A Sepharose Fast Flow (PAS) (GE Healthcare) overnight at 4°C. For FancM immunodepletion, 4 volumes of FancM (1 mg mL⁻¹) or an equivalent amount of rabbit IgG purified from non-immunized rabbit serum (Sigma) were incubated with 1 volume of Protein A Sepharose Fast Flow (PAS) (GE Healthcare) overnight at 4°C. For SPRTN immunodepletion, 4 volumes of SPRTN serum was incubated with 1 volume of Protein A Sepharose Fast Flow (PAS) (GE Healthcare) overnight at 4°C. For SPRTN immunodepletion, 4 volumes of SPRTN serum was incubated with 1 volume of Protein A Sepharose Fast Flow (PAS) (GE Healthcare) overnight at 4°C. For mock depletion, 4 volumes of preimmune serum from matched rabbit, was used. One volume of antibody-conjugated Sepharose was then added to 5 volumes of precleared HSS or NPE and incubated for 1 hour at 4°C. The HSS or NPE was collected and incubated two more times with antibody-conjugated Sepharose for a total of three rounds of depletion. The depleted HSS or NPE was collected and used immediately for DNA replication, as described above.

Protein expression and purification

M.Hpall-His₆, Lacl-biotin, and Lacl-His₆ were expressed and purified as previously described (Duxin et al., 2014). Lysine methylation of M.Hpall was carried out as described (Larsen et al., 2018). Xenopus RTEL1 open reading frame with an N-terminal GST tag separated by a 3C cleavage site was cloned into pFastBac1 (Thermo Fisher Scientific) (pJLS101) using custom gene synthesis from Integrated DNA Technologies (IDT). The RTEL1 sequence was confirmed by Sanger sequencing. Mutants of RTEL1 were created by around-the-horn site-directed mutagenesis, and mutations were confirmed by Sanger sequencing. The GST-RTEL1 Baculoviruses were made using the Bac-to-Bac system (Thermo Fisher Scientific) according to the manufacturer's protocols. GST-RTEL1 and mutants were expressed in 3 L suspension cultures of Sf9 cells (Thermo Fisher Scientific) by infection with RTEL1 baculovirus for 36-48 hr. Sf9 cells were collected via centrifugation and washed with 1XPBS and subsequently pelleted by centrifugation and flash frozen. Cell pellets were thawed and resuspended in an equal volume of 2X Lysis Buffer (100 nM HEPES pH7.5, 1 M NaO₂Ac, 20% sucrose, 0.2% IGEPAL, 4 mM DTT, 2X Roche EDTA-free Complete protease inhibitor cocktail), 1X Lysis Buffer (50 mM HEPES pH7.5, 500 mM NaO₂Ac, 10% sucrose, 0.1% IGEPAL, 2 mM DTT, 1X Roche EDTA-free Complete protease inhibitor cocktail) to the weight of the cell pellet. Cells were lysed by two rounds of sonication, followed by addition of ammonium sulfate (4M stock) to 200 mM final concentration and 45 µl/ml Polymin P (10% stock) and stirred at 4°C for 10 minutes. Lysate was cleared by ultracentrifugation at 25,000 rpm in a Beckman Ti45 rotor for 1 hour. The supernatant was subjected to ammonium sulfate precipitation using 0.2 g/ml ammonium sulfate. Proteins were pelleted by ultracentrifugation at 25,000 rpm in a Beckman Ti45 rotor for 1 hour. The supernatant was discarded and protein pellets were resuspended in 50 mL Wash Buffer A₅₀₀ (25 mM HEPES pH7.5, 500 mM NaO₂Ac, 10% sucrose, 0.01% IGEPAL, 2 mM DTT, 1X Roche EDTA-free Complete protease inhibitor cocktails). The resuspended pellet was incubated for 2 hours with 300 µl of Glutatione Sepharose 4B (GE) at 4°C. Following incubation, resin was first washed with 20 mL of Wash Buffer A₅₀₀ and then with 10 mL of Wash Buffer A₂₀₀ (25 mM HEPES pH7.5, 200 mM NaO₂Ac, 10% sucrose, 0.01% IGEPAL, 2 mM DTT, 1X Roche EDTA-free Complete protease inhibitor cocktails). Proteins were eluted from the resin with Elution Buffer E200 (25 mM HEPES pH7.5, 200 mM NaO2Ac, 10% sucrose, 0.005% IGEPAL, 2 mM DTT, 20 mM L-glutathione reduced, pH adjusted to 8.0). Fractions were pooled and dialyzed against Dialysis Buffer (25 mM HEPES pH7.5, 200 mM NaO₂Ac, 10% sucrose, 0.005% IGEPAL, 2 mM DTT) with addition of HRV 3C protease (Thermo Fisher) at 4°C for 4 hr. Aliquots of RTEL1 were flash frozen and stored at -80°C.

Plasmid pull-down

The plasmid pull-down assay was performed as described (Budzowska et al., 2015). Briefly, streptavidin-coupled magnetic beads (Invitrogen; 10 μ I per pull-down) were washed three times with 50 mM Tris (pH 7.5), 150 mM NaCl, 1 mM EDTA pH 8, 0.02% Tween-20. Biotinylated Lacl was added to the beads (4 pmol per 10 μ I beads) and incubated at room temperature for 40 min. The beads were then washed four times with Pull-down Buffer (10 mM HEPES (pH 7.7), 50 mM KCl, 2.5 mM MgCl₂, 250 mM sucrose, 0.25 mg/ml BSA, 0.02% Tween-20) and resuspended in 40 μ I of the same buffer. The bead suspension was stored on ice until needed. At the indicated times, 4.0 μ I samples of the replication reaction were withdrawn and gently mixed with Lacl-coated streptavidin Dynabeads. The suspension was immediately placed on a rotating wheel and incubated for 30 min at 4°C. The beads and associated proteins were isolated by centrifugation through a sucrose cushion (10 mM HEPES pH 7.7, 2.5 mM MgCl₂, 50 mM KCl, 0.5 M sucrose, 0.02% Tween), then washed once with Pull-down Buffer. All residual buffer was removed, and the beads were resuspended in 20 μ I of 2X LaemmIi sample buffer. Equal volumes of the protein samples were blotted with the indicated antibodies.

DPC pull-down

The DPC pull-down assay is a high-stringency plasmid pull-down protocol to specifically isolate DNA crosslinked Hpall from extract (Larsen et al., 2018). Briefly, Streptavidin-coupled magnetic beads (Dynabeads M-280, Invitrogen; 10 µL per pull-down) were washed

twice with Wash buffer #1 (50 mM Tris pH 7.5, 150 mM NaCl, 1 mM EDTA pH 8, 0.02% Tween-20). Subsequently, biotinylated Lacl was added to the washed beads (1 pmol per 10 μ L of beads) and rotated at room temperature for 40 min. The Lacl-coated beads were then washed four times with DPC pull-down buffer (20 mM Tris pH 7.5, 150 mM NaCl, 2 mM EDTA pH 8, 0.5% IPEGAL-CA630) and stored on ice. Samples (4.5 μ l) from DNA replication or gap filling reactions were withdrawn and stopped in 300 μ L of DPC pull-down buffer on ice. After the final time point, 10 μ L of Lacl-coated streptavidin Dynabeads were added to each sample and allowed to bind for 60 min at 4°C with rotation. The beads were subsequently washed three times with DPC pull-down buffer and then twice with Benzonase buffer (20mM Tris pH 7.5, 150mM NaCl, 2mM MgCl2, 0.02% Tween-20) before being resuspended in 12 μ L Benzonase buffer containing 1 μ L Benzonase (Novagen) and 1 μ l streptavidin-coupled magnetic beads not coated with Lacl (binds any free biotinylated Lacl). Samples were incubated for 1hr at 37°C to allow for DNA digestion and DPC elution, after which the beads were pelleted and the supernatant M.Hpall eluate was mixed with 2X Laemmli sample buffer for subsequent immunoblotting analysis.

Expressing, and purifying recombinant GINS

Codon-optimized *Xenopus laevis* cDNAs encoding all four GINS subunits (synthesized by IDT) were cloned into a single expression plasmid (pGC128) using the MultiBac system (Trowitzsch et al., 2010). A sequence encoding a 10-aa linker, the LPETG tag for sortase labeling, and a His6 tag was added to the C terminus of the Psf3 subunit (GGGGSGGGGS-LPETG-HHHHHH). The bacmid encoding the GINS complex was obtained by electroporating pGC128 into DH10EMBacY (or DH10Bac) electro-competent cells and purified using ZR BAC DNA miniprep kit (Zymo Research Cat# D4048).

Baculovirus encoding GINS was amplified in three stages (P1, P2, and P3) in Sf9 cells (Expression Systems Cat# 94-001S) and GINS expression levels were monitored via western blots. 500mL of Tni cell culture (Expression Systems Cat# 94-002S) at a density of 2-3 million/mL was infected with 5-10 mL P3 baculovirus (MOI > 1). Cells harvested 48 hr post-infection were pelleted at 500 x g for 15min and re-suspended in a final volume of 50mL in GINS Lysis Buffer (GLB) containing 20mM Tris-HCl pH8.0, 5% glycerol, 500mM NaCl, 20mM Imidazole, 1mM DTT, 1mM PMSF, EDTA-free cOmplete protease inhibitor cocktail (Roche Cat # 11873580001). Cells were lysed by sonication on ice and the insoluble fraction was pelleted via centrifugation for 1hr at 30,000 g at 4C. The clarified lysate was incubated with 0.5mL NiNTA resin (QIAGEN Cat# 30410) for 1hr at 4C on a rotating wheel. The resin was washed 5 times with 10mL of GLB in a disposable column. The protein was eluted in 5 rounds with 500uL/each of GLB + 250 mM Imidazole; the elutions were pooled and desalted using a PD10 column (GE Healthcare Cat 17-0851-01) into 20mM Tris-HCl pH7.5, 5% glycerol, 100mM NaCl, 1mM DTT. Recombinant GINS was further purified on a MonoQ column connected to an AKTA Pure FPLC with a 100-1000mM NaCl gradient in 20mM Tris-HCl pH7.5, 5% glycerol, 1mM DTT buffer (the GINS complex eluted at ~500mM NaCl). The eluted protein was desalted using PD10 columns or dialysis into 20mM Tris-HCl pH7.5, 5% glycerol, 150mM NaCl, 1mM DTT; concentrated to ~2mg/mL, frozen in liquid nitrogen, and stored at ~80C. A pellet from 0.5L of Tni cell culture yielded ~5mg of purified GINS.

Purification of Hpall for sortase labeling

A sequence encoding a 10-aa linker, the LPETG tag for sortase labeling, and a His6 tag was added to the C terminus of Hpall (pGC220) and expressed in T7 Express cells (NEB Cat# C2566I) (Duxin et al., 2014). LPETG-tagged Hpall was affinity purified on NiNTA resin, and further purified on a MonoQ column connected to an AKTA Pure FPLC with a 100-500mM KCl gradient in 20mM HEPES pH8.2, 10% glycerol, 0.02% IGEPAL-630, 1mM DTT (Hpall eluted at ~300mM KCl). The fractions containing Hpall were pooled and dialyzed overnight into 20mM HEPES pH8.0, 100mM KCl, 10% glycerol, 0.02% IGEPAL-630, 1mM DTT; concentrated to ~2-3 mg/mL; and stored at -80C. A portion of LPETG-tagged Hpall was methylated as described above, then fluorescently labeled using sortase.

Fluorescent labeling of GINS and Hpall

Sortase-tagged proteins were conjugated to a short peptide labeled with a fluorophore (AF647 or AF568). 1mg of lyophilized GGGGYKCK peptide (synthesized by New England Peptide) was dissolved in 1mL degassed reaction buffer (20mM HEPES pH 7.5, 0.1mM EDTA, 10mM TCEP) and incubated at room temperature for 15 min. 1mg of AF647-maleimide (Thermo Fisher Cat# A20347) or 1mg of AF568-maleimide (Thermo Fisher Cat# 20341) was dissolved in 40 μ L degassed DMSO and added to the dissolved peptide, then incubated on a rotating wheel at room temperature for 3 hr. The reaction was quenched with 10mM DTT. The labeled peptide was separated from unlabeled peptide and un-reacted fluorophore on a MonoQ column with a 10mM-1000mM ammonium carbonate pH 7.8-8.0 gradient (pH adjusted using 50% acetic acid). Fractions containing the labeled peptide were pooled and lyophilized, then dissolved in 50 μ L 20 mM HEPES pH7.5 and stored at -80C.

Conjugation reaction: 2 nanomoles of GINS or 5 nanomoles of Hpall or meHpall were reacted overnight at 4°C on a rotating wheel with 20x molar excess of GGGGYKCK^{AF647} or GGGGYKCK^{AF568} in the presence of 1/10x molar excess of sortase enzyme (purified as previously described (Chen et al., 2011)), supplemented with 100 μ L 4x sortase buffer, in a 400 μ L total volume. 4x Sortase Buffer consists of 80mM HEPES pH 7.5, 600mM NaCl, 40% glycerol, 4mM DTT, and 20mM CaCl₂. After incubation, the reaction mixture was supplemented with NaCl to a final concentration of 500mM and imidazole to a final concentration of 10mM, and incubated with 40 μ L NiNTA resin for 1hr at 4C. NiNTA trapped unlabeled protein and the His6-tagged sortase whereas labeled protein

and excess free peptide remained in the supernatant. Labeled protein was purified and desalted by gel filtration on a Superdex200 column in 20mM Tris-HCl pH7.5, 150mM NaCl, 10% glycerol, and 1mM DTT; then concentrated to 1mg/mL and stored at -80C for up to 1 year.

Preparing substrate with fluorescent DPCs

The DNA substrate was cloned and assembled by tandem recombineering using the MultiBac system (Trowitzsch et al., 2010) as a 30.6 kb plasmid (pGC261) and grown in DH10β cells (NEB Cat# C3019I). The plasmid contains two sites for Hpall conjugation flanked by Nt.BbvCl sites such that a fluorinated oligo can be annealed and ligated into the backbone. 100-150 μg (at ~1000 ng/uL) of pGC261 plasmid was purified using a Midi Prep kit (QIAGEN) from 250mL of bacterial culture. Supercoiled plasmid was nicked with Nt.BbvCI (NEB Cat# R0632S) and the enzyme was heat inactivated. 20-fold molar excess of modified oligo (TCAGCATCCGGTAGCTACT CAATCCGGTACC, the highlighted nucleotide is 5-fluoro-deoxycytidine, synthesized by BioSynthesis) was annealed to the nicked backbone then ligated overnight. The DNA was dialyzed in 10mM TrisHCl pH8.0, then digested with the single cutter NotI-HF (NEB Cat# R3189S) in NEB2 buffer and the enzyme was heat-inactivated. The 3' DNA termini were biotinylated by incorporating biotindCTP (Thermo Fisher Cat# 19518018) and biotin-dGTP (Perkin Elmer Cat# NEL541001EA) using Klenow Fragment 3'→5' exo-(NEB Cat# M0212S). The DNA was then purified by electrophoresis on a 0.6% agarose 1xTAE gel pre-stained with SYBR Safe (Thermo Fisher Cat# S33102), the 30.6kb linear DNA band was excised and electro-eluted in 1xTAE Buffer. The DNA was butanol-extracted several times until the volume was reduced 2-3 fold, then phenol-chloroform extracted 3 times to remove any agarose traces, followed by chloroform extraction. The DNA was ethanol-precipitated and eluted in EB (QIAGEN), and dialyzed in 10mM Tris-HCI H8.0 to remove trace contaminants. Purified DNA was conjugated to 100-fold molar excess of Hpall^{AF568} or meHpall^{AF568} as described above. The reaction was run on 0.6% agarose 1xTAE gel pre-stained with SYBR Safe, the linear DNA band was excised, and the DNA was electro-eluted in 1xTAE. The eluate could be stored in the dark at 4C for a few days or at -80C for at least a few months.

Single-molecule KEHRMIT assay

Immunodepletion of endogenous GINS: For each experiment endogenous GINS was immunodepleted in 3 rounds from 50 µL NPE and in 2 rounds from 70 µL HSS. For each round of NPE depletion 10 µL Protein A Sepharose resin (PAS, GE Healthcare) was incubated with 50 µg of affinity purified anti-GINS antibody at 4C overnight. For each round of HSS depletion, 14 µL PAS resin was incubated with 70 µg of antibody. The resin was pelleted in a swing-bucket rotor for 30sec at 2500 g and washed twice with 1xTBS, once with 1xELB sucrose, twice with 1x ELB sucrose + 0.5M NaCl, and twice with 1xELB sucrose. Immuno-depletions were performed on a rotating wheel in the cold room for 45 min/each.

Flow Cell Assembly and DNA Tethering: Flow cells were assembled using 25x75x1mm slides (VWR Cat# 48300-025), double-sided tape (Grace Bio-Labs Cat# 620001), and PEG-ylated coverslips, as previously described (Yardimci et al., 2012a). 24x60mm No. 1.5 coverslips (VWR Cat# 16004-312) were PEG-ylated using a mixture of 10% Biotin-PEG-SVA and 90% M-PEG-SVA MW5000 (Laysan Bio). Linear DNA biotinylated at both ends (λ DNA or the 30.6kb DNA substrate with dual DPC lesions) was stretched to ~90% of its contour length and immobilized onto the coverslip (Yardimci et al., 2012a). Immobilized DNA was stained with 200 nM SYTOX Green (Thermo Fisher Cat# S7020) and imaged using 488-nm laser excitation (details in the Microscopy section below). 20-50 fields of view (FOVs) near the flow cell inlet were selected and subsequently imaged during the replication experiment every 0.5-3min. The DNA stain was removed by washing the flow cell with 1xELB sucrose before any egg extracts were introduced.

Replication Reaction Mixture Assembly: After immuno-depleting HSS and NPE, the following mixtures were assembled. The ATP regeneration system (ARS) was assembled on ice by mixing 5 μ L 0.2M ATP (Sigma Cat# A-5394) with 10 μ L 1M Phosphocreatine (Sigma Cat# P-6502) and 0.5 μ L 5 mg/mL Creatine Phosphokinase (Sigma Cat# C-3755). Δ HSS-NPE Mix was assembled at room temperature by mixing 31 μ L GINS-depleted NPE with 31 μ L GINS-depleted HSS, and incubated at room temperature for 5min to inactivate Cdt1. The Licensing Mix was assembled by supplementing 20 μ L of GINS-depleted HSS with 2 μ L of 30-bp long dsDNA carrier (300ng/ μ L stock) (Loveland et al., 2012; Yardimci et al., 2012a) and 0.7 μ L ARS, then incubated at room temperature for 5min. The Replication Initiation Mix was assembled by adding 3 μ L pBlueScript carrier plasmid DNA (200ng/ μ L stock) (Lebofsky et al., 2011), 1 μ L ARS, 4 μ L 1xELB Sucrose, and 1 μ L rGINS^{AF647} (0.3-1.0 mg/mL stock) to 18 μ L of Δ HSS-NPE Mix, then incubated at room temperature for 5min. The Replication Elongation Mix was assembled by supplementing 40 μ L of Δ HSS-NPE Mix, twich 7 μ L pBlueScript carrier plasmid DNA (200ng/ μ L stock), 2 μ L ARS, 11 μ L 1xELB sucrose, and, optionally 1-2 μ L of Fen1^{mKikGR} (100 μ M stock), and incubated at room-temperature for 5min.

DNA Replication and Imaging: After the DNA was tethered as described above, replication origins were licensed by flowing 15 μ L of Licensing Mix into the flow cell at 10 μ L/min, and incubating for 2-5min. Replication origins were fired by flowing 20 μ L of Replication Initiation Mix at 10 μ L/min, and incubating for 2-5min. Excess free rGINS^{AF647} was washed away by flowing 55 μ L of Replication Elongation Mix at 10 μ L/min. 2-3 minutes after starting the wash AF568 (for DPC experiments) or mKikGR (for λ DNA experiments) and AF647 were imaged every 0.5-3min for 1-3hrs. Since the PCA/PCD and glucose oxidase/catalase oxygen scavenging systems inhibited DNA replication (data not shown),oxygen scavengers were omitted and stroboscopic imaging was employed to limit photobleaching.

Microscopy: Single-molecule data was collected at the Nikon Imaging Center at Harvard Medical School on a Nikon Ti motorized inverted microscope ("Tobias") equipped with CFI Apochromat TIRF NA1.49 oil immersion objective with 100X magnification, the Perfect Focus System for maintenance of focus over time, a Nikon TIRF illuminator, an Agilent MLC400B laser launch with

405 nm (20mW), 488 nm (50mW), 561 nm (50mW), and 647 nm (125mW) (power measured at fiber optic). Images were acquired using a ZT405/488/561/647rpc dichroic (Chroma) with an Andor DU-897 EM-CCD camera controlled by NIS Elements software.

SYTOX Green fluorescence was excited with the 488 nm laser (< 0.1mW measured at the objective) for 100ms and collected with an ET525/50 m emission filter (Chroma). AF568 fluorescence was excited with the 561 nm laser (0.5-1.0mW) for 100 ms and collected with an ET600/50 m emission filter (Chroma). AF647 fluorescence was exited with the 647 nm laser (0.5-1.0mW) for 100 ms and collected with an ET700/75 m emission filter (Chroma). Fen1^{mKikGR} was photo-switched with the 405 nm laser (0.5-1.0 mW) for 100 ms and collected with an ET700/75 m emission filter (Chroma). Fen1^{mKikGR} was photo-switched with the 405 nm laser (0.5-1.0 mW) for 100 ms and collected with an ET600/50 m emission filter (Chroma). (Loveland et al., 2012).

For time-lapse experiments, images were collected every 0.5-3 min (depending on the experiment) at 20-50 different fields of view (FOVs). Multiple FOV positions were collected using a Prior Proscan II linear-encoded motorized stage. Movie frames for each FOV were aligned in ImageJ using the Image Stabilizer Plugin.

KEHRMIT data analysis

A typical KEHRMIT experiment generated a movie containing several fields of view (FOV), and each FOV initially contained ~100 double-tethered DNA molecules of the correct size (48.5kb \ DNA or 30.6kb DPC substrate extended to ~90% of their respective contour lengths). A significant portion of DNA molecules ruptured when egg extract mixtures were introduced into the flow cell. DNA molecules that were tethered too close to each other aggregated in egg extracts and were omitted from analysis. Molecules where multiple replication origins fired close to each other were also discarded. Only replication origins that gave rise to two labeled CMG molecules were included in the analysis. Active CMG molecules were detected by generating a maximum-intensity z-projection of the CMG^{AF647} channel where translocating CMG molecules appeared as a contiguous bright trail. Regions of interest (ROIs) corresponding to active helicases were hand-selected, automatically cropped, rotated, and individual frames were stacked to generate kymograms. The fluorescent spot image corresponding to individual CMG molecules was fitted by a 2D Gaussian and its position was determined to sub-pixel accuracy. Translocation velocity was computed by linear fitting of CMG position versus time to a straight line. Processivity was measured as the distance traveled by CMG from the beginning of the experiment until the AF647 signal disappeared due to photobleaching, DNA tether rupture, or CMG reaching the end of the DNA template. When analyzing data from aphidicolin uncoupling experiments, it was taken into account that at a low force the length of RPA-bound ssDNA is essentially the same as that of dsDNA (Lewis et al., 2017). In experiments with the 30.6kb DNA-DPC substrate, only DNA molecules containing two labeled DPCs at the expected distance were included in the analysis. Furthermore, DPC^{Lead} and DPC^{Lag} encounters were classified according to the location of the replication initiation event relative to the two lesions as illustrated in Figure 6B. CMG pauses at the DPC were identified as segments of the helicase trajectory where the CMG position was within 1px (160 nm) of the DPC. Unambiguous DPC bypass events (BID) were defined as CMG encounters with a DPC^{Lead} or DPC^{Lag} where both approach and departure could be unambiguously detected (at least 3 time points for each) and the corresponding velocities could be reliably measured. DD+ events were defined as CMG encounters with a DPC^{Lead} or DPC^{Lag} where the DPC^{AF568} signal disappeared but the CMG^{AF647} signal persisted, and CMG traveled at least 1px from the original site of the DPC. Conversely, DD- events were defined as CMG encounters with a DPC^{Lead} or DPC^{Lag} where the DPC^{AF568} signal disappeared, the CMG^{AF647} signal persisted, but CMG traveled less than 1px from the site of the DPC. B+M events were defined as CMG-DPC^{Lag} encounters where both CMG^{AF647} and DPC^{AF568} signals persisted and both CMG and DPC traveled more than 3px from the initial DPC^{Lag} position. Since data was acquired at 1min/frame, pause durations and DPC lifetimes are reported as integers in increments of 1min. Error-bars for mean or median values corresponding to distributions were estimated as 95% confidence intervals via bootstrapping.

KEHRMIT assay strengths and limitations

The KEHRMIT assay offers a few key advantages over fiber analysis - the most common assay used to measure the average speed of replication forks, and PhADE - a real-time assay for imaging the growth of nascent DNA tracts (PhotoActivation, Diffusion and Excitation). (i) KEHRMIT directly monitors CMG dynamics as opposed to DNA synthesis. (ii) KEHRMIT provides temporal information that reveals changes in fork speed or transient helicase pausing. (iii) KEHRMIT has a higher spatial resolution than either DNA combing or PhADE because the position of the CMG helicase on DNA can be determined with sub-pixel accuracy. However, KEHRMIT is incompatible with simultaneous imaging of SYTOX-stained DNA because at the high protein concentration present in Xenopus egg extracts cross-links SYTOX-stained DNA to the surface of PEG-functionalized glass coverslips. Instead, the DNA molecules can be stained and imaged after they are immobilized, and de-stained prior to adding egg extracts to the flow cell. Optionally, DNA can also be SYTOX-stained and imaged after replication in egg extracts and stringent washing with SDS-containing buffer (Yardimci et al., 2012a), but some double-tethered DNA molecules break during replication and subsequently cannot be visualized using SYTOX. In the labeling scheme presented here, dual-color CMG^{AF647} and DPC^{AF568} imaging is incompatible with simultaneous PhADE imaging of Fen1^{mKikGR} bound to nascent DNA (Loveland et al., 2012) because the absorbance/emission spectra of AF568 and the photo-switched form of mKikGR overlap. Consequently, the DPCAF568 signal could not be distinguished from that of Fen1^{mKikGR}. In addition, the use of Fen1^{mKikGR} at very high concentration (1-3 µM) to image nascent DNA synthesis by PhADE precludes the concomitant use of any green fluorophores (such as AF488) because of overlapping absorbance/emission spectra with the green form of mKikGR.

QUANTIFICATION AND STATISTICAL ANALYSIS

All ensemble experiments were performed three or more times, except for Figure 3D which was performed twice. Replication and sequencing gels were quantified in Multi-Gauge software from Fujifilm, the values from independent experiments were averaged and plotted in Microsoft Excel with error bars representing the standard deviation (the relevant statistical details are also listed in the figure legends).

All single-molecules experiments were performed two or more times. Single molecule data was acquired using NIS Elements software from Nikon, pre-processed in ImageJ using the Image Stabilized plugin from Kang Li at Carnegie Melon University, and quantified in MATLAB software from Mathworks. All beeswarm plots represent data from individual molecules as dots (the number of molecules, N, is listed in the figure or figure legends), the mean value is depicted as a horizontal blue line, and the 95% Confidence Interval (CI) for the mean value is illustrated as a gray box. The 95% Confidence Interval for the mean was used to determine if two quantities were significantly different. The standard MATLAB implementation of bootstrapping was used to estimate the 95% CI for the mean value of measurements extracted from single-molecule experiments, as specified in the figure legends.

Supplemental Figures



Figure S1. Disappearance of the CMG Footprint from a Stable DPC, Related to Figure 1

(A) Schematic showing trapping of M.Hpall on a plasmid containing a fluorinated M.Hpall recognition site. (B) Schematic of the two replication-coupled DPC proteolysis pathways in *Xenopus* egg extracts (Larsen et al., 2018). (C) Mock-depleted and SPRTN-depleted egg extracts were blotted using SPRTN and Mcm6 (legend continued on next page)

⁽loading control) antibodies. (D) pmeDPC^{2xLead} was replicated in the indicated egg extracts. At different times, plasmid was recovered under stringent conditions, the DNA digested, and the released proteins subjected to immunoblot analysis with M.Hpall antibody. (E) In Figure 1C, stabilization of the DPC led to the appearance of stalling products from +1 to +12 (lanes 7-12 and 19-24). To determine whether these products depend on TLS, pmeDPC^{Lead} was replicated in the indicated extracts supplemented with $[\alpha - {}^{32}P]$ -dATP. Samples were processed and analyzed as in Figure 1C. This analysis showed that the +1 to +12 products were eliminated by REV1 depletion. These products were also not observed during replication of a normal DPC in SPRTN-depleted extracts containing MG262 (data not shown), suggesting they are specific to DPC methylation. (F) Alternative models to explain the disappearance of the CMG footprint at stable DPCs: CMG dissociation (i) or CMG bypass (ii). (G) Model of CMG dynamics on pmeDPC^{2xLead}. During replication of pmeDPC^{2xLead}, both converging forks encounter a meDPC on the leading strand template. After CMGs bypass the two-leading strand DPCs, CMGs converge, as seen during replication termination, whereupon they are ubiquitylated (probably by CRL2^{Lrr1}) (Dewar et al., 2017), and unloaded by p97. (H) pDPC^{Lead} was replicated in undepleted extract containing $[\alpha - {}^{32}P]$ dATP and supplemented with buffer or 100 µM Cdc7i (PHA767491) at zero or ten minutes after NPE extract addition, which initiates replication, and analyzed as in Figure 1C. The fact that DPC bypass was unaffected when Cdc7i was added immediately after forks reached the DPC argues that new origin firing is not required for DPC bypass. (I) A serial dilution of mock-depleted egg extract was analyzed alongside FancM-depleted egg extracts via blotting with FancM antibody. (J) pDPC^{2xLead} was replicated in the indicated extracts supplemented with [a-32P]-dATP. At different times, DNA was recovered and nicked with Nb. Bsml, separated on a denaturing polyacrylamide gel, and visualized by autoradiography. FANCM depletion inhibited FANCD2 ubiquitylation, demonstrating that FANCM was functionally depleted (data not shown). (K) Upper panel: $pDPC^{2xLead}$ was replicated in the indicated extract containing [α -32P]-dATP and supplemented with either buffer or 66.67 µM ATRi (ETP-46464). Samples were processed and analyzed as in (J). Lower panel: same extracts without addition of $[\alpha^{-32}P]$ -dATP were replicated and reaction samples were separated by SDS-PAGE and blotted with the indicated antibodies.

Figure S2. Models for the Interaction of a 5' to 3' DNA Helicase with Different DPC-Containing DNA Templates, Related to Figure 2 Proposed action of a 5' to 3' helicase (pink) on four different substrates (i-iv). Since there are only 4 bp between the DPC attachment site and the bubble in pmeDPC^{Lag/Lead-Bubble}, arrival of CMG should melt the duplex surrounding the DPC (iv).

Cell

Figure S3. RTEL1 Is Required for CMG Bypass but Not in the Presence of a Converging Replication Fork, Related to Figure 3

(A) Egg extracts contain multiple RTEL1 isoforms, as seen in lane 1 and Figure 3B, lane 1. To distinguish them, we depleted egg extract with an antibody raised against an N-terminal fragment of RTEL1 (α -RTEL1-N, lane 2; used in all experiments except lane 3 of this panel) or an antibody raised against a C-terminal peptide (α -RTEL1-C, lane 3), and blotted with α -RTEL1-N or ORC2 (loading control) antibodies. Unlike α -RTEL1-N, α -RTEL1-C depleted only the largest RTEL1 isoform, consistent with the presence of multiple isoforms (as seen in mice) (Ding et al., 2004), only the largest of which has the C-terminal extension against which the antibody was raised. Depletion of extracts with α -RTEL1-C antibody (lane 3) had no effect on CMG bypass (data not shown), demonstrating that the shorter isoforms were sufficient to perform this function. (B) Incorporation of [α --³²P]-dATP during pmeDPC^{Lead} replication in mock-depleted or RTEL1-depleted extract was quantified and graphed. The mean of n = 5 experiments is graphed. Error bars represent the standard deviation. (C) Coomassie blue-stained SDS-PAGE of purified RTEL1 wild-type and RTEL1-K48R. The RTEL1 and co-purifying GST-tag bands are indicated. (D) pmeDPC^{Lead}, pmeDPC^{Lead}, and pmeDPC^{Lag/Lead}-Bubble were replicated in the presence of [α --³²P]-dATP in the indicated extracts and analyzed as in Figure 31. The mean of n = 3 experiments is graphed. Error bars represent the standard deviation. (E) Quantification of CMG bypass in Figure 3D. (F) DNA samples from Figure 3E were digested with the single-cutter AfIIII, separated on a non-denaturing agarose gel, visualized by autoradiography, and linear products were quantified. The mean of n = 3 experiments is graphed. Error bars represent the standard deviation.

Figure S4. Role of RTEL1 in DPC Proteolysis, Related to Figure 4

(A) pDPC^{2xLead} was replicated in the indicated extracts. At different times, plasmid-associated proteins were recovered under non-stringent conditions and blotted with the indicated antibodies. (B) Extracts used in Figure 4E were supplemented with $[\alpha - {}^{32}P]$ -dATP and used to replicate pDPC^{2xLead}. DNA samples were separated on a native agarose gel and subjected to autoradiography. The gel shows that in the absence of RTEL1, replication products accumulated as slow mobility intermediates instead of undergoing dissolution into monomeric open circular (OC) and supercoiled (SC) plasmids, reflecting a defect in CMG bypass and lack of DNA unwinding of the 165 bp that separates the DPCs (see cartoon). (C) pDPC^{ssDNA} was incubated directly in the indicated extracts without first licensing in HSS, which prevents replication initiation due to the high concentration of Geminin in NPE (Arias and Walter, 2005). Chromatin was isolated by the stringent pull-down procedure described in Figure 4A and analyzed for M.Hpall levels. RTEL1 depletion had no effect on the rate of M.Hpall destruction. (D) Same as in (C) except that M.Hpall conjugated to the gapped plasmid was methylated to inhibit the proteasome pathway. Short and long (larger excerpt) exposures of the same blot are shown. The result shows that in the setting of ssDNA, the SPRTN-specific product was generated at the same rate whether or not RTEL1 was present

Figure S5. KEHRMIT Assay Validation, Related to Figure 5

(A-B) pmeDPC^{2xLead} was replicated in the indicated extracts containing [³²P]-dATP and supplemented with buffer or rGINS^{AF647}. At the indicated times, DNA was recovered, resolved on a native agarose gel (A) or a sequencing gel (B), and visualized by autoradiography. Endogenous GINS depletion abolished and recombinant GINS^{AF647} fully rescued DNA replication (A). rGINS^{AF647} also supported loss of the CMG footprint (B). (C) The efficiency of DPC bypass by CMG from panel B and two additional experiments (n = 3 total experiments) was quantified as described in Figure 2B and graphed. (D) A serial dilution of mock-depleted and GINS-depleted NPE was supplemented with buffer, 300 nM (1x) or 600 nM (2x) AF647-labeled rGINS, or 300 nM (1x) unlabeled rGINS, and analyzed by western blotting with antibodies against the Psf1 subunit of GINS. (E) Integrated fluorescence intensity traces for individual CMG molecules traveling along λ DNA illustrated that AF647 photobleached in a single step. (F) Kymogram of a replication bubble (as in Figure 5C) illustrating that the replication fork continues to travel (blue) after AF647 (green) photobleaching (arrow).

Figure S6. Detailed Analysis of Individual DPC-CMG Encounters, Related to Figure 6

(A) Example of automated analysis of a DPC^{Lead} bypass event. (Left) Kymogram of the CMG-DPC^{Lead} encounter. (Middle) CMG and DPC positions were determined by fitting the centroid of the corresponding fluorescent signal. (Right) Plot of CMG-DPC distance. Approach (green), pause (red), and departure (blue) phases were detected by an automated algorithm (See STAR Methods). (B) Examples of BID, DD+, DD-, CD, and DT events. (C) Photobleaching kinetics for DPC^{AF568} and GINS^{AF647} (N = 200 molecules were analyzed in each case). T_{1/2} denotes the half-life extracted from the exponential decay fit (solid lines) to the raw data (dots). The experiment was conducted in egg extract with DPC^{AF568} and GINS^{AF647} non-specifically bound to the glass coverslip using the same laser power and frame rate as experiments described in Figure 6. These data suggest that AF647 photobleaching is a major contributor to CD events. Although AF568 photobleaching contributes to DD+/– events for stabilized DPCs, it plays only a minor role in DD+/– events for degradable DPCs. (D) Quantification of three different classes of CMG-DPC^{Lag} encounters from n = 2 independent experiments: (i) Bypass of intact DPC (BID), representing unambiguous bypass events;

(ii) CMG bypassed the DPC position followed by DPC mobilization due to sister chromatid separation (B+M). (iii) DPC disappeared first, followed by CMG departure from the pause site (DD+). These are probably bypass events where the DPC was degraded before CMG departure. N represents the number of molecules. (E) Two examples of B+M events at meDPC^{Lag} in SPRTN-depleted extract. (F) Model of how B+M events result from sister chromatid separation when the outward-moving CMG reaches the end of the DNA substrate. (G) Histograms of meDPC lifetime in mock-depleted extract in the absence of replication (N = 199 molecules) (+geminin control to block origin firing) versus meDPC^{Lead} in SPRTN-depleted extracts (N = 151 molecules) and nonmethylated DPC^{Lead} in mock-depleted extract (N = 183 molecules), both in the presence of replication. In all cases, DPC lifetime was measured from the beginning of the experiment. The peak at 90-100 min corresponds to DPC^{AF568} molecules that survived the entire 100-min experiment. Blue lines and gray boxes correspond to the mean and the 95% CI. Given that the DPC lifetime in reactions with active proteolysis (bottom panel) is significantly shorter than the DPC lifetime reactions where both SPRTN and the proteasome are active is largely due to replication-coupled proteolysis and not AF568 photobleaching. (H) Replication origins firing in extracts supplemented with vehicle (left) or 150 µM aphidicolin (right). CMG and nascent DNA signals are shown in green and blue respectively (same as Figure 5C). The absence of blue signal in the +Aph. experiment confirms the inhibition of DNA synthesis. (I) Beeswarm plot of CMG processivity in the presence of aphidiccilin. Blue lines and gray boxes correspond to the mean and the 95% CI. N represents the number of molecules. (J) Beeswarm plot of the time needed to bypass methylated DPC^{Lead} in SPRTN-depleted extract. We replication origins firing in extracts using a set of the inhibition of DNA synthesis. (I) Beeswarm plot of the time needed to bypas

Figure S7. Possible Mechanisms of CMG Bypass, Related to Figure 6

(A) pLacO₃₂ was pre-incubated with LacR and replicated in the indicated extracts containing $[\alpha - {}^{32}P]$ -dATP and supplemented with DMSO or MG262. At the indicated times, DNA was recovered, digested with the single cutter XmnI, resolved by native agarose gel electrophoresis, and visualized by autoradiography. Accumulation of the linear species was quantified, the mean of n = 3 experiments is graphed. Error bars represent the standard deviation. Immunodepletion of SPRTN and addition of proteasome inhibitor had no effect on the conversion of X-shaped intermediates to fully replicated daughter molecules, indicating that LacR proteolysis was not required for replication fork progression through the array. (B) Bypass model in which CMG denatures the DPC and threads the resulting polypeptide chains through the central channel of the MCM2-7. (C) Bypass model in which the MCM2-7 ring transiently opens, possibly due to dissociation of GINS and/or CDC45 (not shown), allowing the helicase to translocate or be towed (not shown) past the DPC. (D) Model for template switching at a DPC. If CMG bypasses the DPC but TLS fails, the leading strand anneals to the nascent lagging strand. After extension of the leading strand (blue arrow), it undergoes dissolution and re-anneals to the leading strand template, thereby bypassing the adduct.